

INTEGRATED ORGANIC AMENDMENTS: A NEW ERA IN VERMICOMPOSTING AND BIOCHAR PRODUCTION DRIVEN BY MICROBIAL ENHANCEMENT OF ORGANIC WASTE

Syda Zahra Haider¹, Ayesha Ali², Sibtain Abbas^{*3}, Mureed Hussain⁴, Samrina Javed⁵
Muhammad Karim Hussain⁶

¹Department of Botany, Ghazi University, Dera Ghazi Khan, 32200, Pakistan

²Department of Botany, University of Agriculture, Faisalabad, 38040, Pakistan

³ Department of Biology, PMAS Arid agriculture university Rawalpindi, 46300, Pakistan.

⁴ Department of Botany, Government College University Faisalabad, 38000, Pakistan.

⁵Department of Chemistry, Kohat University of Science and Technology, Kohat. 26000 Pakistan

⁶Department of Chemical Engineering, University of Engineering & Technology Peshawar, 25000, Pakistan

¹zahrabotanist@gmail.com, ²aansaayesha@gmail.com, ³sibtainabbas1241@gmail.com

⁴mureedmajeed0341@gmail.com, ⁵samrinajaved666@gmail.com, ⁶karimhussain0333@gmail.com,

DOI: <https://doi.org/10.5281/zenodo.17722143>

Keywords

Integrated organic amendments; vermicomposting; biochar; microbial communities; organic waste management; soil fertility; carbon sequestration; sustainable agriculture

Article History

Received: 01 October 2025

Accepted: 10 November 2025

Published: 26 November 2025

Copyright @Author

Corresponding Author: *

Sibtain Abbas

Abstract

The increasing depositional levels of organic waste in agricultural and industrial sources are a great threat to the environment, and its proper management requires sustainable approaches. The possible potential of integrated strategies, especially how vermicomposting and biochar production can be combined to transform these organic residues into useful soil amendments, is also pointed at in this review. Vermicomposting utilizes earthworms and microbes to create compost rich in nutrients that increases soil fertility and soil structure. Biochar is a by-product of biomass pyrolysis, which is a better water retention medium and nutrient availability and long-term carbon storage. These can be applied synergistically: vermicompost is highly microbial and full of nutrient cycling, and biochar contains pores microbiological habitat, which allows the colonization of microorganisms and conservation of nutrients. These not only improves the productivity of the soil but also helps in the rehabilitation of the polluted soils as well as reducing greenhouse gases. Organic matter decomposition and growth promotion of the plant can also be further enhanced by including effective consortia of microbes. Finally, the review states the primary requirements of future research like process optimization, microbial inoculation techniques, and field-based measurements to reinforce the agronomic and environmental application of this form of integrated organic amendments.

INTRODUCTION

World population is rapidly increasing and probable to reach in between 8.5 to 12 billion by 2100. More than 75% of the population is expected to reside in

in the African and Asian countries (Lam et al., 2025). The overpopulation leads to a state of food insecurity that induced the evolution of resource-

exhaustive agriculture causing irreparable environmental damages. With growing pressure on global agricultural systems, the efficient management of organic waste and improvement of soil fertility have become urgent priorities (Chojnacka & Baltrusaitis, 2025). Traditional composting methods, while beneficial, have limitations in nutrient retention and microbial diversity. The integration of vermicomposting and biochar each with distinct properties augmented by microbial consortia, offers a promising solution. This review examines the synergistic potential of these components in promoting sustainable agriculture and environmental remediation and to increase soil fertility for efficient crop production to fulfill the need of ever expanding population. This review, in addition to prominence the effect of vermicompost on soil quality and crop development has evaluated and concentrated the current literature on the progress of research intended at improving the nutrient content of vermicompost by optimizing the factors occupied in the vermicomposting process.

Organic Waste as a Resource

Types of Organic Waste

Organic waste composed of biodegradable materials that were once part of a living organism and go through rapid decomposition. Organic waste has many types and includes Food waste, comprising uneaten, raw or expired food and scraps from households, restaurants, and food processing industries that also constitutes a considerable portion of municipal solid waste. Its disposal contributes to methane emissions and landfill depletion (Dey et al., 2025). However, when properly managed and processed food waste can be converted into valuable resources and organic product. Technologies such as anaerobic digestion and composting offer methods to harness the energy potential of food waste while concurrently reducing its environmental impact (Aryanfar et al., 2025). Yard waste also considered as organic waste that encompasses organic materials like leaves, grass clippings, and branches consequential from landscaping activities. While commonly composted, this can also be utilized for biogas production or biochar generation. By diverting yard waste from landfills, we can alleviate the release of greenhouse

gasses and create cleaner energy alternatives to protect the environment (Dwivedi et al., 2025).

Furthermore, branches, leaves, and other organic materials generated from logging and forestry operations cooperatively form forestry waste. This waste can also be composted and utilized for biomass energy production or may employed in mushroom cultivation. Converting forestry waste into valuable resources and products not only reduces environmental impacts within the forestry sector but also generates opportunities for sustainable livelihoods as well as economic growth (Kumar et al., 2021).


Moreover, animal waste from livestock, poultry and pets commonly known as manure has high nutrient and organic matter content. Through composting, biogas production and fertilizer generation, it can be changed into valuable resources. Harnessing the power of manure not only reduce organic waste and aids waste management as well as reduces greenhouse gas emissions but also enhances soil fertility and promotes sustainable agricultural practices for increasing crop production (Lazaridou et al., 2021).

Agricultural production not only provides basic protection for humanity but also produces a huge quantity of organic materials waste. The situation is getting more and more severe nowadays due to the rapid population explosion and economic growth which has led to an increase in agricultural waste production capacity. Recent studies revealed that approximately 38 billion metric tons of organic wastes are produced all over the world due to agricultural waste (khan et al., 2024). Moreover, to meet the demands of a rising population, livestock and crop production has increased significantly which has further led to the generation of agricultural waste. Human behavior, consumption rate, and population explosion are the generally proposed factors responsible for this dramatic explosion of agricultural waste production. As wastes materials are always considered to be either unusable or disposable and either their burning or deposition has always been the result. Burning and deposition in turn results in numerous environmental problems. Burning pollutes atmosphere while land disposal of organic waste materials may directly or indirectly amend the heavy metal status of the soil by affecting metal solubility and dissociation kinetics

(Oyedeki et al., 2024). In order to deal with this challenging area, various treatment methods and practices have been formulated and applied by

countries all over the world. Figure show different types of organic waste.

TYPES OF ORGANIC WASTE



Challenges & Opportunities

The resource utilization of these organic wastes is an important means to solve the problem especially for Pakistan, one of the countries with the largest agricultural production and ever expanding population. It is expected that the annual output of straw waste is as high as 900 million tons out of which about 30% is not utilized. The annual output of livestock and poultry manure is up to 3.8 billion tons with a comprehensive utilization rate of less than 60% (He et al., 2025). Despite the common wastes such as crop residues and animal manure diversification of agricultural production in Pakistan also produced a diversity of other organic wastes such as biogas residue. These different types of agricultural waste are not properly used that resulting in a series of environmental pollution problems as well. In the meantime, these agricultural organic wastes are rich in organic matter and a variety of nutrients such as nitrogen, phosphorus, and potassium. Therefore, agricultural waste, when used as agricultural organic material, will play an significant role not only in improving the quality of agricultural products but also in improving soil quality and also replacing chemical fertilizers. The heavy use of chemical fertilizers such as pesticides and other non-organic substances, as well as the waste of agricultural organic materials not only

exacerbates the fossil energy and other nonrenewable resource inputs but also caused the environmental pollution and greenhouse gas emissions. Therefore, through the development of recycling agriculture, a huge amount of agricultural waste resources can be “turned into treasure”, which is an important path to cope with energy saving and emission reduction in agriculture (Mehdizadeh et al., 2025).

There are a many of benefits to returning organic materials to the field. The application of various organic materials, such as crop residues and manure, has been noted to improve soil fertility also increase soil organic matter content and even suppress N₂O emissions under some conditions. At the same time, returning organic materials from straw and poultry manure to soil can also increase crop yield. A meta-analysis from previous study showed that returning straw to the field increased the average yield of rice, wheat, and maize yields by 5.04%, 8.09%, and 8.71% respectively (Chang et al., 2023). Furthermore, returning agricultural wastes can to some extent replace the application of chemical fertilizer. World’s agricultural production has been facing challenges from the conflicts between population explosion and arable land limitation. The extensive use of chemical nitrogen fertilizers has caused environmental concerns such as greenhouse gas emissions and water eutrophication. Some studies have reported that organic material returned

to the field replaces 70% of chemical fertilizers and outcome in higher yields than inorganic fertilizers alone because the mixing of manure with chemical fertilizers not only provides essential nutrients but also extends the duration of fertilizer efficiency. In addition, the nutrients in organic materials are released more gradually than chemical fertilizers, which potentially raise the nitrogen recovery rate of the crop and allow for a fuller application of the fertilizer. However, research on organic matter substitution and nutrient utilization is currently inadequate (Bhatt & Kukal 2015).

Returning organic materials to the field also brings some worries. Several recent studies have revealed the influences of diverse organic matters on N₂O emissions, but there are inconsistencies in their results. For example, Wu et al. (2019) observed that straw return increased N₂O emissions in the rice-wheat system. While Li et al. (2020) showed that straw incorporation suppressed N₂O emissions in the wheat growing season. Methane is produced during the anaerobic decomposition of organic matter in landfills. Methane has been responsible for approximately 30% of global warming since the pre-industrial era till now (Zheng et al., 2025). Microbial activity in anaerobic conditions breaks down organic compounds results in producing biogas that is a mixture primarily of methane (50–70%) and carbon dioxide (30–50%) also with trace amounts of other gases (Jia et al., 2025). Landfills and dumps together account for about 11% of global methane emissions that's making them a major anthropogenic source of this persuasive greenhouse gas. Food waste is a particularly significant contributor in this regard. Even though it comprises about 24% of the municipal solid waste (MSW) but it is responsible for an estimated 58% of annual landfill methane emissions. This disproportionate impact is due to the rapid decay rate of food waste which often results in methane formation and emission before landfill gas collection systems can capture it. Even in landfills equipped with biogas recovery systems but fugitive methane emissions persist due to inefficiencies in gas collection and inconsistent degradation rates (Feng et al., 2025). Moreover, Leachate is the liquid byproduct created as water percolates through landfill waste composed of dissolving and suspending organic and inorganic substances. Landfills with a

high proportion of organic waste produce leachate rich in organic pollutants that are characterized by high biochemical oxygen demand and chemical oxygen demand. The composition of landfill leachate is complex and variable and typically including dissolved organic matter such as carbohydrates, fats, proteins, persistent organic pollutants, Inorganic macro-components like salts, ammonia, chlorides, sulfates, heavy metals for example arsenic, cadmium, chromium, lead, mercury and Xenobiotic compounds such as pesticides, pharmaceuticals, industrial chemicals. Leachate pose a significant environmental risk If not properly managed. It can seep into groundwater and surface water and contaminating aquatic ecosystems that threatening biodiversity and posing serious public health risks through the food and water supply chain. The age of a landfill affects leachate composition such as young landfills generate leachate with higher organic content, while older landfills have a tendency to have increased concentrations of inorganic pollutants and heavy metals. Consequently, methane emissions contribute to climate change while on the other hand leachate contamination threatens water quality and ecosystem health (Meegoda et al., 2025).


Managing strategies of organic waste

Managing organic waste involves a wide range of approaches and technologies that intend to extract its potential value while minimizing its human and environmental impact while extracting its potential value. The following are some methods to manage organic waste. Composting is a natural process that turns organic waste into nutrient-rich soil. The process involves piling up organic waste and allowing it to decompose under controlled environmental conditions. Composting can be done at home by means of a compost bin or pile or it can be done on a larger scale at a composting facility. The other process of managing organic waste is anaerobic digestion that breaks down organic waste in the lack of oxygen results in producing biogas and fertilizer (Izzotti et al 2025). This process is usually used to treat sewage sludge agricultural waste as well as food waste. Moreover, Bioreactor technology also involves the use of microorganisms to break down organic waste in a controlled and optimum environmental condition. However, it is important

to make certain that the organic waste is appropriately treated to eliminate pathogens and other harmful substances before its application to land (Meegoda et al., 2025).

However, these processes can have negative population and environmental impacts and its compulsory to be carefully managed to minimize their impact. The other most beneficial and

alternative method of managing organic waste is vermicomposting that is a type of composting that uses various form of worms to break down organic waste into useful product. The worms consume the organic matter and produce a nutrient-rich fertilizer named as vermicompost (Singh et al., 2020).



Vermicomposting: Mechanisms and Benefits

Vermicomposting is the decomposition of organic household and other waste that can be used in combination with inexpensive, space-saving models to improve soil fertility and crop production. Similarly, Waste items obtained from agricultural industry that has not been reused. These agricultural wastes are highly biodegradable and making them a major issue in municipal landfills. These agricultural wastes consist of straw, leaves, twigs, stubbles, and large quantities of grasses and weeds have all been produced during production and cultivation of crops and agriculture (Ahmed et al., 2025).

By functioning together, earthworms and aerobic microorganisms stabilize organic waste during the vermicomposting process. Vermicomposting is a low-cost and environmentally friendly method for

dealing with agricultural waste product. One of the most common earthworm species utilized in vermicomposting is *Eisenia foetida*. Outcome from physicochemical analyses have indicated that in contrast to compost and other agricultural wastes, vermicomposting reduces total organic carbon (TOC) and the carbon-nitrogen (C/N) ratio on the other hand increasing nitrogen-phosphorus-potassium (NPK) content that helps to increase soil fertility and crop production. Additionally, it can be used to increase agricultural yields by eliminating harmful organisms in the environment (biocontrol), enhancing soil water retention, and manufacturing plant growth regulators (Mohite et al., 2024). Vermicomposts can extensively influence the growth and productivity of plants and crops as well due to their micro and macro elements, vitamins, enzymes and hormones. Vermicomposts contain nutrients such as nitrates, exchangeable phosphorus, soluble

potassium, calcium, and magnesium in plant available forms and have large particular surface area that provides many microsites for microbial activity and for the strong retention of nutrients (Shaikh & Shaikh 2025). Uptake of minerals and nitrogen (N), phosphorus (P), potassium (K) and magnesium (Mg) by rice plant was highest when fertilizer was applied in combination with vermicompost. Nitrogen uptake by ridge gourd was higher when the fertilizer mix contained 50% vermicompost. Apart from providing mineralogical nutrients, it also contributes to the biological fertility by adding valuable microbes to soil. Mucus, excreted through the earthworm's digestive canal and also stimulates antagonism and competition between various microbial populations ensuing in the production of some antibiotics and hormone-like biochemicals, boosting plant growth (Gogoi et al., 2025). Chatterjee et al., (2025) reported the nutrient status of vermicomposts with organic carbon 9.15-17.98%, total nitrogen 0.5-1.5%, available phosphorus 0.1-0.3%, available potassium 0.15%, calcium and magnesium 22.70-70 mg/100 g, copper 2-9.3 ppm, zinc 5.7-11.5 ppm and available sulphur 128-548 ppm that is very beneficial for crop production.

Vermicomposts are used as alternative potting media due to their low-cost, excellent nutrient status and physiochemical characters. Considerable improvements in plant growth recorded after amending soils with vermicomposts have been attributed to the physico-chemical and biological properties of vermicomposts that increases the macropore space ranging from 50-500 µm, resulting in improved air-water relationship in the soil, favorably affecting plant growth. It also reduces the proportion of water-soluble chemical, which cause possible environmental contamination. Previous raspberry proves that vermicompost has beneficial buffering capability and ameliorate the damage caused by excess of nutrients which may otherwise cause phytotoxicity. During vermicomposting the heavy metals forms complex, aggregates with humic acids and other polymerized organic fractions resulting in lower availability of heavy metals to the plant, which are otherwise phytotoxic and effects the normal functioning of plants (Ibrahim et al., 2016).

Challenges and Limitations of Vermicomposting

Despite its number of benefits, vermicomposting also faces several challenges and limitations. First of all it is time-consuming process that takes several weeks or even months for the several organic matter to completely convert into vermicompost. This may make vermicompost less practical for people who need a quicker turnaround and results. The time required for complete decomposition can vary depending on the forms of organic material used and environmental conditions. Vermicompost is temperature sensitive process as Earthworms used in it are sensitive to extreme temperatures. High or low temperatures can adversely affect their activity and as well as the efficiency of the composting process. Moreover, vermicomposting needs frequent monitoring and maintenance to guarantee that the earthworms and composting process are in ideal circumstances for better and effective results. This includes monitoring of environmental conditions like moisture content, temperature, and pH. This make it less effective methods.

Furthermore, setting up a vermiculture system may needs an initial investment in infrastructure and ongoing management to guarantee the health of the earthworms and the quality of the compost. if someone does not manage the composting pile properly a strong, unpleasant odor can arise from the breakdown process. This may be an issue for individuals or communities who live near to the composting facility. Farmers lack the necessary knowledge to implement vermicomposting systems effectively as successful vermiculture requires a certain level of expertise and knowledge.

Due to its limitation other process may needs to explore to manage agricultural as well as other waste. The other useful process for managing organic waste may be known as bichar whose details are explained below.

Biochar: Properties and Soil Amendment Potential

Biochar is a material rich in carbon that outcome from the pyrolysis process of any waste material of various types such as wood, fruit shells, distiller's grains, press cake from vegetable oil and juice production, rice and nut hulls, residues of plants, manure, industrial and municipal waste, sewage sludge, farming, and fermentation residuals. Biochar

remains in the soils for thousands of years and it is therefore considered more resistant to decomposition than organic matter, acting as an important long-term carbon sink. Biochar typically contains 50 to 80 percent carbon, depending on the substrate and processing parameters. Biochar properties can also be effected by the characteristics and properties of the processing parameters and equipment. The environmental conditions like control of temperature, heating rates, holding time, off-gasses, use of steam or other carrier gases, and physical handling systems can all impact the quality, structure, composition and behavior of the final product. For example, biochar from rapid and fast pyrolysis reactors tends to be very finely grained, as a result of the physical and thermal processes used by the equipment (Talwar et al., 2025).

Biochar is formed from organic material that either grows naturally or is grown as a result of agricultural activity. It may be manure from farm animals, or is waste material such as cardboard, waste wood material, or other organic material that could not be recycled or reused. The carbon content of the given material was absorbed and obtained from the atmosphere through the process of photosynthesis. So, first and foremost Biochar in its basic form is a carbon neutral material. It can even be consider of as "carbon negative" if its end use involves a long lifetime in the form of elemental carbon rather than carbon dioxide (Parvari et al., 2025).

Biochar is mainly used in soils to enhance soil aeration, decrease soil emissions of greenhouse gases, diminish nutrient leaching, cut soil acidity, and potentially boost the water content of coarse soils. Biochar application may amplify soil fertility and agricultural productivity as well. However, when applied excessively or made from feedstock unsuitable for the soil type, biochar soil amendments also have the potential for negative effects, including harming soil biota, dropping available water content, varying soil pH, and escalating salinity (Rasaq et al., 2024).

As a result of all these variables, biochar's properties can vary extensively. It can be hydrophobic or hydrophilic and can have changeable amounts of surface area, diverse reactivities, different strength characteristics, and unstable electrical properties. In addition, further processing of the biochar often in

the form of either "activation" or "graphitization" can modify those properties even more, giving it valuable characteristics for different end uses and application. Because of this, it is important to know what "kind" of biochar you are either producing or seeking to purchase. Since biochar can be formed from such a wide array of low-value materials, there can be little to no competition for the diversity of materials from which it can be created. In fact, production of biochar can often help decrease amounts of unwanted waste by consumption of materials that do not otherwise have a marketable value. Furthermore, since biochar can be formed from invasive plants such as weed trees and other similar materials, it is possible to use biochar as a management tool to advance the local ecosystem through strategic harvest of overgrown or undesirable species (Ren et al., 2025).

Biochar is prepared using a thermal process that has been baked at an elevated temperature in an oven-like device called a "retort." At its heart, this is not an overly complex process, and most of us have created biochar at some point by, for example, leaving bread in the toaster too long is also form of bichar. Biochar construct can be either a dry process called roasting or a wet process named as hydrothermal processing. Usually, roasting is less expensive, but hydrothermal processing can be worth considering, especially if your biomass feedstock has very elevated moisture content. During the thermal process, much of the dry mass is lost from the biomass, and yields can range from 90% down to 10%. The temperature of the process can range from 200 to over 1000°C, and the duration of the treatment may vary from a fraction of a second to several hours, depending on the desired result and outcomes of the final products. In all cases, the amount of oxygen available to the feedstock must be significantly limited - otherwise, the feedstock has a inclination to burst into flames. Occasionally, elevated pressure is also used to affect the thermal processing of the biomass especially when wet processing is used for biochar production (Ayaz et al., 2025).

Another way to think about biochar production is that biochar can be manufactured either as a primary product or as a byproduct of another process. In terms of primary or dedicated biochar creation, a large number of equipment can be used, ranging

from something as simple as a 55-gallon steel drum to something as sophisticated as a continuous flow, computer-controlled industrial retort designed to manufacture tons of biochar per hour. The elevated temperature used to convert biomass into biochar can come from an externally applied source of heat that is often generated by burning a portion of the feedstock. Biochar as a byproduct is char that is left after some other material is created from biomass and parent substrate. The most likely source of byproduct biochar is from either fast pyrolysis results in creation of biocrude oil or from gasification for synthetic natural gas production. In both cases, the biochar is a small portion of the original mass of the feedstock (Meng et al., 2025).

Biochar can be used as a soil amendment to potentially enhance yields, advance soil structure, raise water holding capacity, reduce soil pentation resistance, enhance pH, raise nutrient absorption and decrease nitrous oxide emissions. When the suitable amount and variety is applied to the soil, it can remediate the overflow of excess water and nutrients while retaining absorption for plants during stress conditions. With the copious pores openings and plenty internal surface area it provides a substrate for microbial growth as well as for anchoring root systems as a "reservoir" to accumulate moisture and nutrients. Choice of the type of biochar is very imperative. A high specific surface area (SSA) biochar apply to soils at high rates can bind up herbicides and pesticides, making them ineffective. One study evaluated that that using a high specific surface area biochar (EUC-800) at high rates exaggerated the efficacy of sulfentrazone herbicide while a low specific surface area biochar at the same rate did not. In some cases, when a high specific surface area biochar was apply at high rates, the herbicides were not effective at all, even at the highest label rates. Since biochar has a half-life of hundreds and possibly thousands of years, using the wrong product or the wrong rate could have long-term effects on the agronomic practices that work and may have drastic effect on yield (Santos et al., 2025).

Biochar can be adapted for used as a remediation agent to confine organic and inorganic contamination in the soil or water. It will combine to heavy metals, toxic chemicals, pesticides, and

nutrients to keep them from translocating in the soil profile. This could be used at abandoned mine sites or on roadside runoff, in tile line effluent, or in stormwater basins to capture pollutants and water-soluble nutrients. This trait can pose a problem if the wrong type of biochar is used, as it can bind up soil-applied herbicides used to control weeds. In other applications, biochar can be used as a filtration medium in much the same manner while as activated carbon. It is useful in filtering both liquids and gases as it absorbs various contaminants as well as odor-produce constituents related to manures and municipal wastewater treatment operations. Biochar has a carbon matrix structured with a high degree of porosity and extensive surface area, permitting it to act as a sorbent and play a crucial role in controlling pollutants in the environment. Biochar can help farmers enhance farm land productivity; concurrently decrease the environmental footprint of farming practices by substituting for chemical pesticides, herbicides and fertilizers. Biochar has recently received greater than before attention due to its wide range of agricultural benefits. The adding up of biochar into farm land increase the amount of organic matter content in the soil, thereby improving soil physicochemical and biological properties. Biochar can positively or negatively affect the soil microbial growth to alter the agricultural environment as well as yield and crop production (Tanwar et al., 2025).

Environmental impact of biochar

The formation of functional groups and adsorption sites both on the surfaces and inner pores of the biochar enhance the soil's CEC. Due to this effect, biochar has been used for absorption of organic and inorganic contaminants or pollutants (Sharma et al., 2025). Moreover, the bioavailability of heavy metals such as lead [Pb] and cadmium [Cd]) can be decreased by immobilizing them in soil using biochar as an amendment which limits their movement through the soil. This role of biochar is the consequence of the negatively charged groups on its surfaces, which seems to raise with time due to the oxidation processes in the soil. Biochar beside with phytoremediation strategies could give a good combination for effectively stabilizing and decontaminating heavy metal-polluted sites. On the

other hand, some researchers have warned against health hazards due to exposure to polycyclic aromatic hydrocarbons. Biochar application to agricultural soils and farmland has been recommended as a strategy to diminish increasing atmospheric carbon dioxide concentrations and alleviate climate change (Zhang et al., 2023). Several studies have reported that biochar application decrease the emissions of soil greenhouse gases but the mechanisms responsible for the effects of biochar on soil GHGs emissions are still unclear and needs further investigation. The biochar feedstock application rates increase nitrous oxide emission. Moreover, the increase in soil methane and CO₂ emissions were also reported after biochar application.

CONCLUSION

The application of biochar in improving soil health, crop productivity, land reclamation and reduce of climate change has been extensively studied over the last decade. This research has shown varied results, from clear improvements of soil health to almost no difference compared with the control, depending on the types of soil, crop, biochar and the environment. Moreover, information is still very limited from large scale field trials for biochar application, probably because of the high costs of biochar in the market. Large-scale production of biochar from all types of organic wastes may help in reducing the costs and its practical uses in field. The greatest potential may be in having very targeted applications to address specific environmental concerns. A proper regulatory mechanism and controlled supervision is required to avoid any possible risks of soil contamination due to potential contaminants present in biochar and also on the global GHGs emissions when producing it, which should be considered when using it as a carbon sequestration strategy.

Microbial Inoculants: Enhancing Functional Efficiency

Microbial inoculants are psychrophilic microorganisms that are applied to soil or plants to enhance nutrient availability and plant growth by improving the functioning of the ecosystem and enhancing fertility of the soil. These inoculants usually entail positive microbial communities inclusive of nitrogen fixers, phosphate solubilizers,

decomposers, and useful fungi including arbuscular mycorrhizal fungi (AMF). Nitrogen fixing bacteria such as *Rhizobium* are essential in the movement of nitrogen in the atmosphere to bio-available forms that can be used by plants thus aiding sustainable food production because it limits the use of chemical fertilizer. Phosphate solubilizing bacteria, like *Pseudomonas* species, enhance the bioavailability of phosphorus by solubilising the phosphate compounds that are insoluble in soils. In the meantime, *Trichoderma* and other decomposers play a role in breaking organic matter down and increasing nutrient cycling and AMF form symbiotic interactions with plant roots to nutrients (especially phosphorus) and make plants healthy (Samantaray et al., 2024; Saeid, 2022).

These microbes usually interact in a synergistic fashion with organic amendments such as biochar and vermicompost that forms a habitat and substrate of microbial colonization. The porous structure of biochar allows protective niches which make the survival and activity of microbes possible, and vermicompost is a high-quality source of nutrient and microbial population. Biochar and vermicompost co-application has also been reported to be effective in improving microbial biomass, enzyme activity, nutrient availability and general microbiological environment in soil resulting in better plant growth and yields. This synergism effect is associated with enhanced nutrient bioavailability, and inhibition of soil pathogens, which develop a favorable atmosphere where crops can grow (Saeid, 2022).

The strategies to enhance the functional effectiveness of organic amendments through the improvement of native or introduced microbial communities are bioaugmentation and biostimulation. Bioaugmentation is the intentional application of specific isolates or consortia of microbes in the soil or organic supplements to enhance degradation, nutrient recycling, or promotion of plant growth. Biostimulation is the process of changing environmental conditions, nutrient supplement or aeration to activate the behavior and proliferation of native beneficial microbes. Such strategies coupled with organic amendments can go a long way in improving microbial colonization, nutrient mobilization and general soil health. Nevertheless, to

achieve successful bioaugmentation the inoculant forms of microbes need to be chosen wisely and be able to effectively colonize and functional in a particular soil condition (Saeid, 2022).

To sum up, microbial inoculants which include nitrogen fixers, phosphate solubilizers, decomposers, and AMF are important in sustainable agriculture. Their synergetic relations with biochar and vermicompost enhance the availability of nutrients and the quality of soils. Biostimulation and bioaugmentation methods also maximize microbial functional performance of the organic amendments, which can provide future opportunities in terms of improving the crop yield and productivity as well as soil health in eco-friendly manners. It utilizes microbial diversity, organic sources, and specific management of the microbes to develop sustainable and productive agroecosystems (Samantaray et al., 2024).

Future opportunities and challenges of microbial inoculants to improve functional efficiency contain various significant points which should be considered in detail so that it will be possible to develop microbial inoculants to be used in agriculture. The issue of standardization is a significant concern, which requires the working out of sound and generally accepted protocols to be used in the preparation of organic amendments and microbial inoculants. Present laws and quality standards on microbial inoculants are diverse in different parts of the world, and much is being done to standardize the requirements. To illustrate, regulatory frameworks of AMF inoculants consist of the definition of viable counts of propagules, composition of products, infectivity potential and safety evaluations to determine efficacy and to avoid ecological hazards. Another task associated with standardization is the labeling and packaging of products and bioassays and molecular characterization performance validation. These standards facilitate reliability, consistency, and confidence of the farmers besides supporting international trade and commercialization (Saeid, 2022).

There is also the issue of scalability especially the scaling up a successful inoculant formulations and small-scale testing to the field scale that is uniformly successful and effective. Field conditions like soil

type, climate, and crop species are variable and may affect individual microbial performance and survival thus complicating large-scale application. Multi-location field trials should be done in the long-term to confirm the efficacy of the inoculants under different environmental conditions and to determine the optimal procedures of application. They should also test microbial persistence, effects on native soil microbiomes, and crop responses in the several growing seasons in field tests (Samantaray et al., 2024).

Farmer adoption is highly dependent on economic viability. The expenses of microbial inoculant manufacturing, quality monitoring, formulation, storage, and distribution should be offset by the actual benefits like the rise in the output, the decrease in the application of chemical fertilizers, and enhanced soil health. Evidence of cost-effectiveness by showing effective impact analysis and life cycle analysis can be used to promote its adoption by farmers, particularly in resource-starved environments. There is further support of acceptance and proper use through outreach, education, and demonstration projects (Saeid, 2022).

The gaps in research in this area are diverse:

- Consistent benefits need to be determined using field trials over a long period, behavior of inoculants should be understood, and environmental safety should be monitored. Mechanistic investigations of microbial interactions, such as colonization behavior, synergistic activities when interacting with organic amendments such as biochar and vermicompost, and competition with native microorganisms are essential in formulations optimization.
- To measure ecological footprints, economic returns, and sustainability benefits of using microbial inoculants in agriculture, life cycle analyses and overall impact have to be conducted (Samantaray et al., 2024).

With concerted research and policy action, it is possible to overcome these challenges, which will lead to practical implementation of microbial inoculants, which will allow sustainable increase in

soil fertility, crop yields, and the state of the environment in general.

Summary:

Incorporation of vermiculite (vermiculite vermiculite) and biochar when used synergistically as part of effective microbial communities may be characterized as a ground-breaking and integrated sustainable waste management and soil health regeneration strategy. Vermicompost, which contains valuable microorganisms, enzymes and growth promoting substances like cytokinins and humic acids promotes nutrient availability, soil structure and the activity of microorganisms. Biochar, in the meantime, acquires stable, porous substrates that enhance physicochemical characteristics of soils like pH, porosity, water retention and bulk density as well as offers a habitat protecting and enhancing the growth of healthy microbial populations. The combination of these amendments forms an enabling environment which improves nutrient cycling, disease suppression, and the general soil fertility, which promotes healthy plant growth and production. It has been shown that biochar combined with vermicompost has synergistic effects that are even more beneficial than the individual use of each and each application yields better crop yields, an improved nutrient retention and a healthier soil microbial ecosystem. The pore structure of biochar enables colonization and growth of microorganisms found in vermicompost which enhances microbial activities including decomposition of organic matter and solubilization of nutrients. This combined system also prevents the bioavailability of the heavy metals and toxicity in the soils, therefore saving the plants, hence sustainable functioning of the agroecosystem. These synergistic systems are very promising to be integrated into the concept of the circular bioeconomy in the future by transforming the agricultural and organic wastes into useful resources that can restore the degraded soils and eliminate the use of chemicals. By developing more resilient agroecosystems to environmental challenges, they help mitigate climate change by monitoring long-term soil carbon storage. The interaction of vermicomposting, biochar, and selected inoculation of microorganisms, therefore, represents a future-oriented approach toward improving soil health,

circuitry, and the sustainability of agriculture at large scale, which makes them the basis of efficient future practices of eco-friendly farming and waste valorization.

REFERENCES

- Ahmad, A., Mahmood, A., Ahmad, S., & Li, W. (2025). Harnessing Earthworms for Sustainable Waste Management: Insights into Vermicomposting. *Water, Air, & Soil Pollution*, 236(13), 840.
- Aryanfar, Y., Garcia Alcaraz, J. L., Keçebaş, A., Fernandez, J. B., Arslan, B., Ilbas, M., ... & Irshad, K. (2025). Transforming food industrial sludge into sustainable resources: innovations in waste management and renewable energy recovery. *Business Strategy and the Environment*, 34(3), 3672-3700.
- Ayaz, M., Muntaha, S. T., Baltrėnaitė-Gedienė, E., & Kriaučiūnienė, Z. (2025). Biochar and carbon-negative technologies: exploring opportunities for climate change mitigation. *Biochar*, 7(1), 17.
- Bhatt, R., & Kukal, S. S. (2015). Soil water dynamics and water productivity of rice-wheat system under different establishment methods. *Ludhiana, Punjab: Punja Agricultural University*.
- Chang, G., Liu, H., Yin, Z., Wang, J., Li, K., & Gao, T. (2023). Agricultural production can be a carbon sink: A case study of Jinchang City. *Sustainability*, 15(17), 12872.
- Chatterjee, R., Debnath, A., & Mishra, S. (2020). Vermicompost and soil health. In *soil health* (pp. 69-88). Cham: Springer International Publishing.
- Chojnacka, K., & Baltrusaitis, J. (2025). Organomineral fertilizers for sustainable agriculture. *Sustainability Science and Technology*, 2(2), 022001.
- Dey, S., Santra, M., Choudhury, M., Ghosh, A. R., & Samanta, P. (2025). Food waste generation and its industrial utilization: An overview. *Environmental Science and Pollution Research*, 32(12), 7493-7512.

- Dwivedi, A., Kumar, U., Singh, P. K., Kumar, I., Singh, P., Mishra, S., ... & Sharma, R. K. (2025). Treatment Methods for Food Waste. In *Resource Recycling and Management of Food Waste* (pp. 49-88). Cham: Springer Nature Switzerland.
- Feng, M., Lin, Y., Liu, J., Ni, X., Cheng, Y., Hu, H. W., ... & He, J. Z. (2025). Chinese milk vetch incorporation inhibits nitrification by suppressing comammox Nitrospira in subtropical paddy soils. *Geoderma*, 461, 117469.
- Gogoi, R., Borah, M. P., Anshu, Sharma, M. R., Barooah, M., & Bora, S. S. (2025). Vermicompost Use in Sustainable Agriculture: Impact on Soil Fertility and Plant Growth. In *Innovative Technologies for Waste Management: Strategies, Challenges and Future Directions* (pp. 427-449). Cham: Springer Nature Switzerland.
- He, D., & Cui, C. (2025). Fermentation of organic wastes for feed protein production: Focus on agricultural residues and industrial by-products tied to agriculture. *Fermentation*, 11(9), 528.
- Ibrahim, M. H., Quaik, S., & Ismail, S. A. (2016). Vermicompost, its applications and derivatives. In *Prospects of organic waste management and the significance of earthworms* (pp. 201-230). Springer, Cham.
- Izzotti, A., Pulliero, A., Khalid, Z., Ferrante, O., Aquilia, E., Sciacca, S., ... & Ferrante, M. (2025). Using Natural Isotopes for the Environmental Tracking of a Controlled Landfill Site for Non-Hazardous Waste in Liguria, Italy. *International Journal of Environmental Research and Public Health*, 22(4), 528.
- Jia, L., Liu, T., Chen, Y., Liu, H., Wu, C., Wang, H., ... & Yang, J. (2025). Comparative study on the application of straw, biochar, dimethylpyrazole phosphate, and polyaspartic acid on nitrogen cycle, soil properties, and yield of potted rice. *Plant and Soil*, 1-17.
- Khan, M. N., Sial, T. A., Ali, A., & Wahid, F. (2024). Impact of agricultural wastes on environment and possible management strategies. In *Frontier studies in soil science* (pp. 79-108). Cham: Springer International Publishing.
- Kumar, A., Adamopoulos, S., Jones, D., & Amiandamhen, S. O. (2021). Forest biomass availability and utilization potential in Sweden: A review. *Waste and Biomass Valorization*, 12(1), 65-80.
- Lam, D. (2025). The Next 2 Billion: Can the World Support 10 Billion People?. *Population and Development Review*, 51(1), 63-102.
- Lazaridou, D. C., Michailidis, A., & Trigkas, M. (2021). Exploring environmental and economic costs and benefits of a forest-based circular economy: A literature review. *Forests*, 12(4), 436.
- Li, N., Ma, X., Xu, H., Feng, Y., Ren, G., Yang, G., ... & Ren, C. (2020). Biochar addition mitigates nitrogen loss induced by straw incorporation and nitrogen fertilizer application. *Soil Use and Management*, 36(4), 751-765.
- Meegoda, J. N., Chande, C., & Bakshi, I. (2025). Biodigesters for sustainable food waste management. *International journal of environmental research and public health*, 22(3), 382.
- Meegoda, J. N., Chande, C., & Bakshi, I. (2025). Biodigesters for sustainable food waste management. *International journal of environmental research and public health*, 22(3), 382.
- Mehdizadeh, M., Omid, A., Matindike, R., Nigussie, Z. G., Ikegwu, T. M., Agu, H. O., ... & Merghany, R. M. (2025). Agri-Waste Valorization: Pathways to Sustainable Bioenergy and Biochemical Innovation. *Circular Economy and Sustainability*, 1-31.
- Meng, Z., Xiong, H., Zhu, C., & Xu, G. (2025). Energy saving and carbon reduction production process of municipal sludge-based biochar: Optimization, simulation, and techno-economic analysis. *Process Safety and Environmental Protection*, 193, 577-588.
- Mohite, D. D., Chavan, S. S., Jadhav, V. S., Kanase, T., Kadam, M. A., & Singh, A. S. (2024). Vermicomposting: a holistic approach for sustainable crop production, nutrient-rich bio fertilizer, and environmental restoration. *Discover Sustainability*, 5(1), 60.

- Oyedeeji, S., Patel, N., Krishnamurthy, R., & Fatoba, P. O. (2024). Agricultural wastes to value-added products: economic and environmental perspectives for waste conversion. *Biowaste to Value-added Products: Economics and Technologies*, 215-248.
- Parvari, E., Mahajan, D., & Hewitt, E. L. (2025). A Review of Biomass Pyrolysis for Production of Fuels: Chemistry, Processing, and Techno-Economic Analysis. *Biomass*, 5(3), 54.
- Rasaq, W. A., Okpala, C. O. R., Igwegbe, C. A., & Białowiec, A. (2024). Navigating pyrolysis implementation—a tutorial review on consideration factors and thermochemical operating methods for biomass conversion. *Materials*, 17(3), 725.
- Ren, S., Yang, S., Chen, H., Wang, L., Liu, M., Wang, G., & Xu, C. (2025). A state-of-the-art review on the utilization of biochar as renewable energy for the sustainable steel industry. *Applied Energy*, 394, 126188.
- Saeid, A. (2022). An insight into microbial inoculants for bioconversion of wastes to bio-organic fertilizers. *Environmental Microbiology Reports*, 14(5), 515-530.
- Samantaray, A., et al. (2024). Advances in microbial based bio-inoculum for amelioration of soil fertility and plant growth. *Frontiers in Microbiology*, 15, 117150.
- Santos, L. O., Silva, P. G. P., Machado, B. R., Silva, I. Q., Zapparoli, M., de Brito, G. M. K., ... & da Rosa, A. P. C. (2025). Thermochemical Conversion of Microalgal Biomass: A Potential Source of Bio-oil, Biochar, and Syngas. In *Microalgae as Promising Source of Commercial Bioproducts: Advances and Emerging Technologies in Bioactive Compounds, Biofuels, Food and Biorefineries* (pp. 59-69). Cham: Springer Nature Switzerland.
- Shaikh, K. M., & Shaikh, R. U. (2025). Vermicompost: role in sustainable agriculture. *Infinite Path Res Across Disciplines*, 125.
- Sharma, B., Ghimire, R., Sapkota, S., Shrestha, P., Brewer, C. E., & Adhikari, S. (2025). Nitrous oxide mitigation potential of biochar derived from agricultural and forest biomass: Effects of feedstock composition and pyrolysis temperature.
- Singh, A., Karmegam, N., Singh, G. S., Bhadauria, T., Chang, S. W., Awasthi, M. K., ... & Ravindran, B. (2020). Earthworms and vermicompost: an eco-friendly approach for repaying nature's debt. *Environmental Geochemistry and Health*, 42(6), 1617-1642.
- Talwar, P., Agudelo, M. A., & Nanda, S. (2025). Pyrolysis Process, Reactors, Products, and Applications: A Review. *Energies*, 18(11), 2979.
- Tanwar, V. S., Singh, S. K., Sharma, J. K., & Patidar, R. (2025). Innovative Approaches to Pretreatment and Modification of Sewage Sludge-Derived Biochar for Resource Recovery: A Short Review. *Journal of Building Material Science | Volume*, 7(03).
- Wu, L., Wang, Q., Shang, Z., Chen, H., Deng, A., Song, Z., ... & Zhang, W. Straw Return Boosts N₂O Emissions Primarily During the Pre-Jointing Stage in Winter Wheat Fields: A Meta-Analysis. Available at SSRN 5392113.
- Zhang, X., Lv, J., Zhang, Y., Li, S., Chen, X., & Sha, Z. (2023). A meta-analysis study on the use of biochar to simultaneously mitigate emissions of reactive nitrogen gases (N₂O and NO) from soils. *Sustainability*, 15(3), 2384.
- Zheng, J., Huang, X., Fan, J., Zhang, F., & Wang, H. (2025). Strategies for nitrogen management to mitigate nitrous oxide and ammonia emissions from maize field while maintaining yields in China: findings from a field experiment and a national meta-analysis. *Journal of the Science of Food and Agriculture*.