Policy Research Journal
ISSN (E): 3006-7030 ISSN (P) : 3006-7022 Volume 3, Issue 12, 2025

VITAMIN D FORTIFICATION IN DAIRY PRODUCTS: EVALUATING POLICY IMPACT AND NUTRITIONAL OUTCOMES IN SUNLIGHT-LIMITED REGIONS

Tayyba Iqbal[*1], Ziporah musoke[2], Saba Anayat Laiba[3]

[*1]*University of central Punjab Department of food and nutritional sciences University college Birmingham*
[2]*Department of food and nutritional sciences University of central Punjab Lahore*
[3]*Department of food and nutritional sciences University of central Punjab Lahore*

[*1]taibaseo13@gmail.com, [2]z.musoke1@gmail.com, [3]sabaanayat2005@gmail.com, [4]anjum7496750@gmail.com

DOI: https://doi.org/10.5281/zenodo.18149976

Keywords
Vitamin D deficiency, Dairy fortification, Nutritional outcomes, Policy impact, Sunlight limitation, public health nutrition

Article History
Received: 03 November 2025
Accepted: 17 December 2025
Published: 31 December 2025

Copyright @Author
Corresponding Author: *
Tayyba Iqbal

Abstract
Vitamin D deficiency is a common world health problem, especially in areas with limited sunlight. The purpose of this review is to investigate the usefulness of vitamin D fortification of dairy products as a sustainable nutritional intervention to curb deficiency and enhance health outcomes. The aim is to determine the current policy on fortification, technology, and nutritional effects in various geographical and demographic environments. The analysis of data based on global studies and policy reports was made to evaluate the efficacy of mandatory and voluntary fortification programs to enhance the levels of serum 25-hydroxyvitamin D. The fortification of dairy products, particularly milk and yogurt, has greatly enhanced the population level of vitamin D status in such countries as Finland, Canada, and Sweden, which has led to a decrease in the incidence of deficiency and related health problems such as rickets, osteoporosis, and immune dysfunctions. Even in most developing nations, such as Pakistan and certain Asian countries, such policy discrepancies, inadequate consumer awareness, and variable fortification are still in existence. The broader effects of fortification programs are hampered by policy issues, such as a lack of regulation, stakeholder or agency coordination, and inadequate monitoring. The review concludes that a unified, evidence-based policy system, regular monitoring, and education of the population are essential to reach the maximum advantages of vitamin D fortification. It suggests international cooperation in order to have fair accessibility to fortified food, particularly to the susceptible in sunlight-deprived regions.

INTRODUCTION

Introduction

Vitamin D or calciferol has taken a new turn in the limelight in recent years, and the scientific studies conducted regarding the subject have been closely examined. Vitamin D is a vitamin that is lipid-soluble and plays a key role in the regulation of phospho-calcium that ensures that calcium denotes are maintained at normal levels and assists in the development of the bone. Vitamin D is a vitamin-like compound that is made up of two derivatives: vitamin D3, aka cholecalciferol, and vitamin D2, also called ergocalciferol, both of which are essential. The two forms of vitamin D are significant in the body since they guard the body against diseases like rickets or bone erosion, high blood pressure, malignancies, and immunodeficiency diseases. It also plays a central role in the anti-infective defense through the effect of anti-inflammatory, immunomodulatory, proapoptotic, and antiangiogenic activities. A molecule known as Gc-globulin or vitamin D binding protein ferries vitamin D in the body, and it is also considered very important to ensure that the body is at homeostasis, as it plays an anti-inflammatory role [1].

The role of vitamin D has increased exponentially over the past few decades, particularly after the scarcity of vitamin D was found to be closely associated with numerous diseases, and moreover, in the world, vitamin D deficiency appears to be very common. Nowadays, the role of vitamin D as a hormone is confirmed in numerous physiological and pathophysiological activities, in relation to various organs and systems of the human body. Despite extensive data revolving around the skeletal effects of the vitamin D hormone, irrespective of age, there are vigorous arguments as to whether the extra-skeletal effects of the vitamin D supplement, and extremely large doses, can be harmful. Most of the researchers are, however, in agreement that patients who have a deficiency (or an insufficiency) of vitamin D must be placed under therapy to enhance their bone health and general health in general. This is particularly the case in high-risk deficient patients, such as older age (particularly the long-term care centers), diabetes, chronic kidney disease (CKD), malabsorption, and others [2,3,4,5].

It is important to maintain adequate levels of 25-hydroxypytoinol D [25(OH)D] (more than 30 ng/mL) of the vitamin in serum in community-dwelling, postmenopausal women though it is not advised that these women take vitamin D supplementation to protect against fractures in primary prevention of the disease rather than preventing the occurrence of incidences of hypocalcemia in case of secondary prevention amongst patients, who already have suffered fragility fractures. When it comes to the field of vitamin D deficiency, the knowledge on this deficiency is dubious and insufficient, though there is rising awareness about the consequences of vitamin D deficiency. The need to address vitamin D deficiency is generally ignored due to high prevalence nowadays, ease of diagnosis, and the fact that the answer to this issue is simple, effective, and very inexpensive [2,3,4,5].

In fact, vitamin D is also a fat-soluble prohormone steroid with endocrine, paracrine, and autocrine activities. Vitamin D is involved in the endocrine activity of maintaining serum calcium. Vitamin D and calcium are normally referred to as one sentence, as they are the two that are supposed to work in collaboration. The one that is supposed to control these levels of calcium in the bloodstream is vitamin D, which constantly permits the assimilation of calcium and phosphate to enter the body via the intestine, or the bones to lose calcium. Further, vitamin D is a facilitating aspect, which, at ideal concentrations of its own effect on calcium absorption, exerts no perceptible effects, although it permits or enables physiologic plasticity to the changing needs of calcium [6,7].

According to the statements, dietary intake tends to provide an insignificant amount of total vitamin D supply, but it is still a substantial source. It is assumed that vitamin D intake, as it is integrated into daily intake of food, mostly assessed by a standard mass-study food-frequency questionnaire and/or 24-hour recall technique, is well below 10 mg /day in many European communities. Vitamin D intake among different age groups in Northern Europe varies between 3-14 4g/day, in Western Europe between 1.5-5 4g/day, in Southern Europe between 1- 3 4g/day, in Eastern Europe between 2-5

4g/day, and in the Middle East between 1-4 4g/day. Moreover, non-Western immigrants have been seen to have low levels of serum 25(OH)D in European countries compared to the indigenous dark-skinned people [45].

Vitamin D Deficiency in Sunlight-Limited Regions

Deficiency of vitamin D has been cited to be a significant public health problem, which is steadily growing at a rate globally. Vitamin D is produced as a result of the synthesis of UV-B in the skin, but the surrounding environments and antiquated lifestyles tend to curtail sunlight exposure. Reduced exposure to the sun, as well as a sedentary lifestyle, tends to cause the further spread of the vitamin D deficiency. In addition, orthopedic complications are rare few with a considerable number, which normally impacts the well-being of patients. Vitamin D deficiency has been linked to adverse effects on mothers, neonates, and children [8]. Sickle cell disease (SCD) and vitamin D deficiency have similar features that include bone complications and bony pains. The impairment of sun exposure throughout the year and possible deficiency of the diet, combined with high nutritional demands caused by SCD, may result in the disruption of the vitamin D status in Canadian SCD children [9].

Vitamin D plays a vital role in bone and mineral metabolism, skin, immune, and vascular functioning. Nevertheless, more than one billion adults and children around the globe have been estimated to have vitamin D deficiency. Vitamin D in human beings comes through cutaneous production, vitamin D- D-fortified diet, and vitamin D supplements, with at least 80 percent consisting of the former. Adequate levels of ultraviolet B(UVB) radiation trigger the transformation of 7 7-dehydrocholesterol (7DHC) to previtamin D3 in skin. Being a significant external influence factor on vitamin D conflation, UVB radiation depends on season, latitude, and rainfall conditions. Further, internal factors associated with vitamin D status have also been found to be dependent on age, skin saturation, vulnerable skin area, as well as time of sun exposure. China is a diverse nation with numerous geographical regions, intensity of sun radiation, good health practices, and style of dressing, hence the vibrant phenomenon of vitamin D deficiency across regions [10,11,12,13]. At higher latitudes, such as those of the Nordic countries (55–72°N), sunlight is insufficient in both quality and quantity to support UVB-induced vitamin D_3 synthesis in the skin for several months each year, a period often referred to as the "vitamin D winter". According to estimates by O'Neill et al., this period lasts about eight months in Tromsø, Norway (69°N), and around six months in Copenhagen, Denmark (55°N). Furthermore, natural dietary sources of vitamin D are limited, primarily including fatty fish, egg yolks, and certain wild mushrooms. Consequently, vitamin D insufficiency, defined as serum 25(OH)D levels below 50 nmol/L, has been commonly observed across the Nordic population [52].

Sickle cell disease (SCD) and vitamin D deficiency share several overlapping characteristics, particularly bone-related complications and pain. Limited sun exposure throughout the year, coupled with insufficient dietary intake and the increased nutritional demands associated with SCD, can significantly affect vitamin D status in children with SCD living in Canada [53]. The main aim of this review is to give a full overview of the scientific knowledge of Vitamin D fortification of dairy products. It tries to understand the biochemical importance of Vitamin D, its importance in bone and general health, and why dairy products are considered efficient delivery methods in fortification. Through this mix and match of data provided in various studies, the review aims at providing an evidence-based overview on how fortification strategies can be designed, implemented, and optimized to solve Vitamin D deficiency in populations with low sun exposure. The other important objective is to examine the current policy frameworks of Vitamin D fortification in different regions critically. This involves the appraisal of the effectiveness of the existing models, their nutritional performance, and overall health effects on the general population, especially in regions that are at low levels of ultraviolet (UV) radiation. The review will also aim at establishing major research and policy gaps that cripple the ability to roll out large-scale fortification programs and their sustainability. The review will identify the gaps and suggest evidence-based solutions to improve the two

fortification strategies, as well as policy coordination to improve the health outcomes of the populations around the world.

Vitamin D and Human Nutrition

Several studies have continually indicated that the amount of vitamin D is still sub-optimal in various regions of the world, mainly as a result of inadequate exposure to the sun. Vitamin D is found in a limited number of foods naturally, good sources of vitamin D 3 being some types of fish (not only fatty fish), yolks of eggs, and organ meat such as liver. In the same countries, there are countries in which certain foods, such as milk are fortified with vitamin D in an attempt to enhance intake. But total intake of food supplies of vitamin D across the world is still low due to the low food supplies globally. The use of vitamin D supplements is rather significant in maintaining normal levels and is widespread in certain groups. However, as in times of inadequate UVB exposure, current dietary guidelines are, in most cases inadequate in order to obtain or sustain optimal serum 25(OH)D levels. There is hence a recommendation that the intake levels be increased to at least 10 mg per day in all ages in cases where sunshine is restricted, but elderly persons might need up to 25 mg per day [54].

Vitamin D exists in the most important forms: ergocalciferol (vitamin$_2$) and cholecalciferol (vitamin D_3). Vitamin D_2 is produced when ergosterol in yeast and fungi is exposed to ultraviolet light, and people get it from natural foods. Vitamin D_3, on the other hand, is formed from 7-dehydrocholesterol, determined in the skin`s keratinocytes when exposed to sunlight. When nutrition D_2 is consumed, its absorption begins off evolved withinside the stomach, wherein the enzyme pepsin breaks down proteins, and is kept in the duodenum, wherein different digestive enzymes release nutrition D from the food. Bile acids then assist in shaping combined micelles containing fat-soluble compounds for emulsification and synthesis. These micelles are absorbed via way of means of enterocytes and transported first via the lymphatic system and which into the bloodstream [55].

Functions of Vitamin D

Vitamin D and bone health:

Extreme loss of vitamin D can cause rickets in toddlers or children, or osteomalacia in adults, however those are uncommon illnesses withinside the majority of evolved nations. Subclinical nutrition D deficiency is greater common, though, and can be associated with osteoporosis and multiplied occurrence of falls or fractures. The technique of bone mineral deposition begins in pregnancy, mainly withinside the 1/3 trimester of pregnancy. A natural, balanced food plan containing adequate quantities of calcium is higher than normal calcium supplements [18].

Vitamin D and the Innate Immune System:

The first is the innate immune system, the immune system which is needed to fight against pathogens the quickest way possible. Vitamin D has been a well-recognized regulator of innate immunity; however, the initial information on this subject has been produced on the therapy of illnesses caused by mycobacteria, including tuberculosis and leprosy; nonetheless, the process behind these effects has been clarified over a recent period [19].

Vitamin D and cardiovascular health:

Data from observational research advocate that low ranges of 25(OH)D can negatively have an effect on CV health. Currently, however, there may be no consensus concerning the premier stage of the nutrition for viable preventive CVD or most cancers benefits. Vitamin D is idea to play a shielding position towards CVD due to its receptor [20].

Vitamin D and Muscle Function:

The extreme depletion of vitamin D is marked by muscle weakness, but whether milder deficiency or insufficiency of vitamin D can accelerate the mass and performance of muscles, or whether it contributes to the likelihood of falling-related injuries and deaths, is not well described. Clinical implications of the age-related decreases in the mass and muscle performance are significant, including frailty, increased risk of falls and fractures, among other injuries that accompany the fall risk and death. Even vitamin D plays a great role, which is evidenced by the fact that the muscles are extremely weak in the

case of frank vitamin D deficiency. The significance of vitamin D deficiency on the muscle is not so certain. The paper is intended to comment on the muscle alterations during old age, their similarities to changes related to the lack of vitamin D, and the influence of vitamin D supplementation on the muscle mass, their balance, and the risk [21].

Vitamin D Fortification in Dairy Products

Concept of Food Fortification

Among the factors that would make any policy of industry-wide fortification important would be the establishment of a regulatory measure of the addition of the nutrient, through a code of practice or otherwise. This would help prevent the addition of excesses to individual foods or to the total food supply, perhaps by limiting the amount that can be added to individual foods and /or the choice of foods that can be fortified. Depending on the age, food preferences, region, or any other related factors, the selection of food vehicle (and form of the vitamin, i.e., D3 vs. D2) would allow one to target certain groups of consumers. Consideration should be given to factors like: consumer acceptability and preference; shelf-life of the fortified foods and stability of the fortificant over an extended period; and relative healthiness of the food vehicle (so that no fortificant is applied to food that is not considered to be healthy) [22].

Supplementation and meals fortification are the various maximum not unusual place techniques to cope with insufficient day by day nutrition D consumption and to save you or control health problems related to nutrition D deficiency. While supplementation packages depend upon elements which include population coverage, person compliance, cost, and regular consumption, meals fortification overcomes a lot of those obstacles through reaching a broader section of the populace greater effectively. An endorsed and secure method to make sure giant effect is to beef up more than one meal automobiles with smaller nutrition D quantities in place of enriching unmarried meals supply with an excessive dose. Commonly fortified staples which includes milk, margarine or oils, wheat, rice, and maize flours, consisting of bread, function green automobiles for nutrition D fortification. This method has already been applied in numerous countries, consisting of the USA, Canada, India, and Finland, and its effectiveness has been always supported through several systematic evaluations and meta-analyses over the last decade [56].

Why Dairy Products Are Ideal Vehicles

Dairy products and milk intake are healthy for all ages, but more so in children and adolescents since the growth of bone mass is active. Not only do milk and dairy products contain highly bioavailability calcium, but vitamin D and proteins are also sources of the nutrient. Milk and dairy products should be avoided or minimized by patients with lactose intolerance or an allergy to cow's milk, as they contain high-availability proteins, vitamin D, calcium, and phosphorus, which are valuable preventive factors against disorders that may occur in the elderly, e.g., osteoporosis. It is important to mention that lactose intolerance is frequently widespread in geriatric patients, and among such patient populations, fermented dairy products (kefir, yoghurt) and lactose-free milks are the most optimal option. It is important to mention that cow milk and plant milk are different products, and it is impossible to discuss plant milk as a completely valuable substitute for cow milk [23]. When we're talking about adding vitamin D to food, picking the right food and the right type of vitamin D is really important. A lot of studies have looked into whether these foods actually do a good job of boosting vitamin D levels in people. But other things, like how the food tastes, how much it costs, and what people think about vitamin D in their food, haven't gotten as much attention [24].

The combination of all of the above traits of milk, which makes it an ideal delivery system of vitamin D, leads to the choice of certain governments to pressure or even to enforce companies that manufacture certain types of food to add vitamin D to a substantial degree to do so, based on sound nutritional arguments. This holds with foods, typically having very low levels of the vitamin, that are regularly and daily consumed, which can be their source, provided they are fortified, of a consistent and nutritionally pertinent supply of the vitamin. Specifically, in Finland (as well as spreadable fats) fortification of milk (and in Sweden, of milk, vegetable drinks aimed at displacing milk: 0.38–0.50

µg/100 g) is highly encouraged (1 µg/100 g in milk, yogurt and sour milk), whereas in Sweden, it is compulsory (below 3% fat) in milk (as well as in vegetable drinks that are to be treated as substitutes to milk: 0.38 0.50 µ Canada (milk: 0.825 -1.125 mg/100 g) has a similar approach [25].

Can higher doses of vitamin D be toxic?

Concerns about vitamin D toxicity persist for healthcare providers and public health organizations. Despite the lack of recent scientific evidence, this remains a significant reason why governments oppose fortifying foods like milk and dairy with vitamin D. Prior to the 1950s, vitamin D was commonly added to various foods as it was seen as a miracle nutrient effective in treating chronic diseases, including tuberculosis and rheumatoid arthritis [52].



Nutritional Outcomes of Dairy Fortification

Similar to other fat-soluble vitamins, vitamin D bioavailability is regulated by three mechanisms: bio accessibility—release of the lipid phase of the food matrix and solubility in mixed micelles; transformation—biochemical and biochemical transformations; and transport and uptake—migration through the mucus layer to the intestinal lumen surface and uptake by epithelial cells. Bio accessibility is likely one of the three phenomena influencing bioavailability; therefore, it is possibly the most important one. The food matrix effect on vitamin D bioavailability is not fully known. Lipids can also boost vitamin D bioavailability in ways including but not restricted to: formation of a hydrophobic phase in which it can be solubilized; reduction of biliary secretion and resultant production of micelles; and the creation of lipids to produce vitamin D in addition to the production of micelles and lipid digestion products of lipids that promote the production of chylomicron synthesis which aid in the transport of vitamin D beyond the enterocyte. Still, they have not yet been backed by a body of in vivo research available showing that adding more fat to meals would boost the bioavailability of vitamin D. Dietary fiber seems to obstruct vitamin D via reducing the creation of a micelle and by increasing the viscosity of the chime, hence restricting the spread of vitamin D carrying micelles to the brush border. Still, the findings of the in vivo investigations presently available are unable to support this conclusion [33].

Health Improvements

Global Vitamin D deficiency is a concern that is relevant to all ages. The elderly is the population who suffer the most, but youngsters and healthy people are also affected; indeed, the normal process of getting vitamin D from the body might not be good enough in people from sunny areas, and this could be mainly due to the changing lifestyles over the past few decades. Consequently, vitamin D deficiency is one of the emerging and widespread problems of public health because vitamin D is the main element that controls and regulates the metabolism of calcium and bone health. Moreover, a lot of new studies have come up with more information that points to its involvement in the functioning of the immune system, and mainly its ability to stop the formation of inflammatory cytokines and to operate in every part of the human body, which helps to lower the occurrence of many diseases that are considered common diseases. In order to help the vitamin D levels in the population, which are not satisfactory due to the limited common diet intakes, food fortification has become a regular practice and one of the most accepted ways to improve the vitamin D status of the population. At the top of the list is cow's milk, which is a food with great importance for a large percentage of the global population and which is very nutrient-dense for a very low price. Moreover, the contemporary methods for milk sanitizing and packaging are such that the consumption of milk can be kept at room temperature for up to 10-12 months, thus making

the supply of this food accessible to an increasing trading area. In such a scenario, milk could be a handy means of giving a daily adequate intake of vitamin D to keep the homeostasis in the blood, thus reducing the risk of insufficiency or deficiency. Even though the natural content of vitamin D in cow's milk is low, i.e., 0.1-1 μg/L in full-fat milk, the vitamin D fortification of milk for consumption has proven effective in enhancing the overall vitamin D status of the population [34].

The most practical way to raise vitamin D levels is by consuming greater amounts since UVB exposure varies by nation and time of year and carries a chance of skin cancer. Adherence to supplements is constrained, as is nutritional intake of dietary vitamin D from food sources naturally high in vitamin D2 and/or D3. Hence, policymakers, scientists, and producers should turn to the most often consumed foods while thinking about vitamin D reinforcement in foods. Using conventional fortification or biofortification methods, meals can be enhanced with vitamin D2 or D3 [35].

Breastfeeding is among the most vital factors for early mortality, growth, nutrition, and maternal health during the first year of life. Along with the United Nations Children's Fund, the World Health Organization recommends that for the first six months of life babies should just be given breast milk. Still, it is well established that breastfed children especially in high-latitude areas run vitamin D deficiency. The major cause is that the vitamin D concentrations in breast milk depend on the mother's vitamin D levels, which are generally low [36]. Moreover, depending on where the babies are housed, their parents' culture, ideas, customs, or other factors, their exposure to sunlight may be quite constrained. Many of the top medical institutions worldwide therefore support vitamin D supplementation for babies, particularly those exclusively breastfed. For example, AAP issued a guideline in 2008 stating that vitamin D supplementation should be started in the first few days of life for both breastfed and partially breastfed newborns at a dose of 400 IU/day and continued until the child is fully weaned to at least 1 L or 1 qt of vitamin D-fortified formula or whole milk per day. Past studies have shown that there are some barriers to vitamin D supplementation in babies; one of those is the doctors' belief that the kids in their neighborhood get enough sunlight throughout the day or that milk has adequate vitamin D, hence making it unnecessary for doctors to write prescriptions. Taking diet D even as pregnant impacts the physical stages of diet D (specially 25(OH)D) in moms and their babies and can additionally have an effect on maternal insulin resistance and fetal growth. In order to strengthen exercise and coverage regarding the extent of diet D for keeping a wholesome pregnancy, there's a want for exquisite randomized medical trials that examine the pregnancy-unique 25(OH)D thresholds and clinically suitable final results studies [36].

Strategically fortified and/or biofortified staple foods will lessen the risk of future deficiency; yet, modelling fortification/biofortification techniques is needed to protect customers from being exposed to excessive quantities of vitamin D. Therefore, we need to know the treatment impact of vitamin D food supplementation/biofortification across population categories including males and women as well as obese and normal weight people so that we can create safe and effective means of boosting dietary intake of vitamin D [37].

Challenges in Sunlight-Limited Regions

Environmental Barriers

The need to deficit vitamin D is nowadays denoted as a global health problem, regardless of daylight supply. This has questioned the fact that the physical demands can be met by the herbal solar publicity and the downstream occasions. Although the solution to the cutting-edge deficiency requires short-term period methods in form of supplementation or fortification, there should be an implementation of a long-term solution to the problem in the coming years. The issue of huge deficiency in diet D is therefore as far as sunlight publicity is concerned the sole viable method to the problem which is mostly applicable in the case of tropical nations. In addition, diet D-unbiased blessings of solar publicity has also been projected and thus, solar publicity can never be modified completely by fortification or supplementation. As an illustration, UV publicity is hypothesized to boost the production of opioid beta-endorphins which are temper elevators and UVA produced nitric oxide can improve heart health2. It

has been associated with higher mortality in a dose-sensitive manner in Sweden, which has been marked by higher solar-in search of behaviour3. Moreover, a range dependency has been identified in the case of atheroma of the carotid artery and cardiovascular mortality3. This leads to the necessity of finding the prospects of a balanced solar publicity in the population corporations that inhabit the tropical countries. The aim of the currently used evaluation was to examine the clinical evidence that was linked to the challenging conditions and possibilities of application of solar publicity as a sustainable approach to combating a deficiency of diet D by reviewing the trials that involved the use of solar publicity as an intervention. The use of synthetic reagents of UVB as an intervention in studies has now been abandoned. Additional factors (environmental and otherwise) such as precursor, pores and skin colour, calcium status, safety, and various micronutrient deficiencies, which influence the intake of diet D during solar exposure, in the process of continuing ecological and vitamins transition have also been reviewed [38].

Cultural and Lifestyle Constraints

Humans are predicted from a physiological perspective to obtain more than 80% of their vitamin D demands through ultraviolet sunrays given their limited dietary intake. Over the last several decades, there have been major behavioral changes: more people are avoiding the sun and working indoors. Thus, a worldwide epidemic has resulted from clinically significant vitamin D deficit. Globally burden of acute and chronic illnesses and healthcare expenditures are exacerbated by vitamin D deficit [39].

Alternative Food Vehicles

Because of their reputation for being more environmentally friendly, healthier, and more ethical, consumers are becoming more and more interested in factory-grounded goods.1 −3 One of the most significant orders of this growing demand are factory-grounded derivatives of bovine milk, akin to oat, almond, coconut, rice, and soy milks. Though they have well-known environmental and moral advantages, there have been some nutritional profile questions. For example, many plant-based milks lack essential micronutrients (e.g., calcium and vitamin D), which could cause long-term health issues if they are not gotten in adequate amounts from other sources. Thus, the nutritive fortress of plant-based milk analogs may be a great technique to prevent any deficiencies in these trace elements. Originally intended to enhance almond milk, vitamin-enriched nano emulsions were created using surfactant (quillaja saponin) and a factory-grounded oil painting (sludge oil painting) [40].

Research Gaps and Future Directions

One area needing more research is the nutritional and policy divide among people with little access to vitamin D-fortified dairy products. Because of its widespread consumption, high nutritional density, and capacity to provide steady fortificant, dairy makes a great carrier for vitamin D enhancement. Still, differences in consumption patterns, fortification levels, and access mean that some demographic groups continue at greater risk of deficit. Although they are based on antiquated and methodologically limited research involving just 155 people, current milk consumption criteria in the United States were developed mostly to boost dietary calcium intake. Moreover, there are several other healthier sources of calcium and the wider cultural and behavioral context of milk intake should be taken into account when assessing its health effects.

Sophisticated understanding of the connection between dairy intake and health outcomes has come from large-scale epidemiological studies. Although these relationships might be confused by healthier life choices among dairy users, the PURE study—with more than 138,000 people from 21 low- and middle-income nations with rather low baseline dairy intake—discovered that greater dairy intake was somewhat linked with lower risk of cardiovascular disease and all-cause mortality. Subsequent Mendelian randomization investigations of almost 1.9 million people utilizing genetic variations associated with lactose metabolism as a substitute for milk intake have been done to reduce such prejudices; these analyses confirmed relationships between greater milk intake and elevated obesity risk as well as identified lower blood cholesterol levels. Taken together, these results show that while milk and dairy products remain excellent and practical

means of vitamin D fortification, long-term, high-volume consumption of cow's milk may not be optimal for all groups. Along with fortified food plans, a balanced approach stressing moderation and dietary variety seems to represent the most health-conscious strategy [59].

Vitamin D is among the vitamins set to soon turn centennial. Study interest and effort have increased dramatically in recent years. PubMed claims that papers including vitamin D in their title increased from one in 1922 to 110 in 1952, 740 in 1982, and 5566 in 2020. Part of the current spike in research interest is the realization that vitamin D contributes to non-skeletal activities and calcium and bone homeostasis, all of which may be impaired by vitamin D deficiency. Ten years ago, or thereabouts, we examined current research demands and information gaps in vitamin D nutrition and public health, with particular emphasis on the prevention of vitamin D deficiency. Focusing on three research needs in particular will enable us to quantify vitamin D deficiency in the population, prevent it, and possibly prevent disease by increasing vitamin D status throughout the range of knowledge gaps. These were to collect data on the distribution of(i) serum 25- hydroxyvitamin D (25(OH) D) attention and(ii) vitamin D inputs and food sources in nationally representative populations with relevant consideration of coitus, life- stage and race, and(iii) development of sustainable food- founded methods to close the difference between current and recommended intake of vitamin D to reduce vitamin D insufficiency [41].

The success of microencapsulation of Vit D in medicinal lead to its operation in food with the following objectives(i) overcomes solubility hedge between vit D and the food matrix(ii) security vitamin D against physicochemical stress similar as humidity, oxidation, pH, temperature, mechanical and so on(iii) assures better bioavailability with controlled and targeted release of reprised vitamin D [43].

Conclusion

The assessment of the vitamin D fortification of dairy products highlights the necessity of vitamin D in remedying the lack of vitamin D in the world, including areas with low access to sunlight. Policy-based interventions in Finland, Canada, and Sweden have shown that the fortification of milk and dairy items has resulted in substantial changes in the level of vitamin D, bone strength, and general wellness in the population. These results accentuate the use of dairy products as the best fortification vehicles, given that they are more consumed, have nutrient synergy with calcium, and can be readily incorporated into existing diets. Although these successes have been achieved, there are still a number of challenges. Lack of organized policies on fortification is still experienced in many developing countries, which leads to continued high levels of deficiency. Poor awareness among the population, poor policy enforcement, and technological factors hinder development. Also, voluntary fortification in some countries adds to the unequal availability of vitamin D in food products. To achieve uniform benefits, it is necessary to strengthen the mandatory fortification policies, strictly control the quality, and encourage the industry to comply.

Fortificant have been made more palatable to consumers with technological advances such as microencapsulation, which enhances the stability and sensory properties of fortificants. Nevertheless, fortification programs should also be successful with the help of constant education of the population aimed at facilitating consumption and addressing the lack of awareness regarding sources of vitamin D. A combination of fortification and supplementation interventions and the diversification of diets can offer a long-term solution to decreasing the prevalence of deficiency in the world. To summarize, fortification of dairy products with vitamin D is an efficient, inexpensive, and sustainable approach to enhancing the health outcomes of the population in areas that have restricted access to sunlight. Policymakers, nutritionists, and food industries should work together to come up with evidence-based guidelines, uniform fortification levels, and permanent monitoring mechanisms in order to bring in their full potential. The benefits of such coordinated efforts will not only be improved population vitamin D status but also make a substantial contribution to the reduction of the burden of deficiency-related diseases on the global population.

Moving forward, the optimization of the level of fortification to combine the efficacy and safety of fortification in the various groups of the population should be considered in future research. Individualized nutrition, based on genetic diversity, food practices, and environment, may enhance the fortification interventions to become more effective. In addition, the incorporation of digital health technologies and population-scale monitoring can add to the collection of data and monitoring of long-term results. International efforts are crucial, especially when it comes to low- and middle-income countries, to provide equal opportunities to access fortified foods. Through further innovation and policy incorporation, the fortification of dairy products with vitamin D can still be developed as one of the pillars of nutritional health and disease prevention in the world.

REFERENCES

Pop, T. L., Sîrbe, C., Benţa, G., Mititelu, A., & Grama, A. (2022). The role of vitamin D and vitamin D binding protein in chronic liver diseases. *International journal of molecular sciences*, *23*(18), 10705.

Dominguez, L. J., Farruggia, M., Veronese, N., & Barbagallo, M. (2021). Vitamin D sources, metabolism, and deficiency: available compounds and guidelines for its treatment. *Metabolites*, *11*(4), 255.

Charoenngam, N., & Holick, M. F. (2020). Immunologic effects of vitamin D on human health and disease. *Nutrients*, *12*(7), 2097.

Holick, M. F. (2007). Vitamin D deficiency. *New England journal of medicine*, *357*(3), 266-281.

Pike, J. W., & Christakos, S. (2017). Biology and mechanisms of action of the vitamin D hormone. *Endocrinology and Metabolism Clinics*, *46*(4), 815-843.

Mostafa, W. Z., & Hegazy, R. A. (2015). Vitamin D and the skin: Focus on a complex relationship: A review. *Journal of advanced research*, *6*(6), 793-804.

Vanchinathan, V., & Lim, H. W. (2012, April). A dermatologist's perspective on vitamin D. In *Mayo Clinic Proceedings* (Vol. 87, No. 4, pp. 372-380). Elsevier.

Elsori, D. H., & Hammoud, M. S. (2018). Vitamin D deficiency in mothers, neonates and children. *The Journal of steroid biochemistry and molecular biology*, *175*, 195-199.

Grégoire-Pelchat, P., Alos, N., Ribault, V., Pastore, Y., Robitaille, N., & Mailhot, G. (2018). Vitamin D intake and status of children with sickle cell disease in Montreal, Canada. *Journal of pediatric hematology/oncology*, *40*(8), e531-e536.

Holick, M. F. (2017). The vitamin D deficiency pandemic: Approaches for diagnosis, treatment and prevention. *Reviews in endocrine and metabolic disorders*, *18*(2), 153-165.

Saraff, V., & Shaw, N. (2016). Sunshine and vitamin D. *Archives of disease in childhood*, *101*(2), 190-192.

Wacker, M., & Holick, M. F. (2013). Sunlight and Vitamin D: A global perspective for health. *Dermato-endocrinology*, *5*(1), 51-108.

Fan, P., Wang, Q., Li, J., Lu, C., Xu, Y., Cao, H., ... & Chen, D. (2021). Poor status of vitamin D: a survey of area with lowest sunlight radiation in Sichuan, China. *Frontiers in Endocrinology*, *12*, 626983.

Moulas, A. N., & Vaiou, M. (2018). Vitamin D fortification of foods and prospective health outcomes. *Journal of biotechnology*, *285*, 91-101.

Institute of Medicine (US), Food and Nutrition Board, & Committee to Review Dietary Reference Intakes for Vitamin D and Calcium. (2011). *Dietary reference intakes for calcium and vitamin D*. National Academies Press.

Calvo, M. S., & Whiting, S. J. (2003). Prevalence fo Vitamin D insufficiency in Canada and the United States: Importance to health status and efficacy of current food fortification and dietary supplement use. *Nutrition reviews*, *61*(3), 107.

Calvo, M. S., Whiting, S. J., & Barton, C. N. (2004). Vitamin D fortification in the United States and Canada: current status and data needs. *The American journal of clinical nutrition*, *80*(6), 1710S-1716S.

Chang, S. W., & Lee, H. C. (2019). Vitamin D and health-The missing vitamin in humans. *Pediatrics & Neonatology*, *60*(3), 237-244.

Sassi, F., Tamone, C., & D'Amelio, P. (2018). Vitamin D: nutrient, hormone, and immunomodulator. *Nutrients*, *10*(11), 1656.

de la Guía-Galipienso, F., Martínez-Ferran, M., Vallecillo, N., Lavie, C. J., Sanchis-Gomar, F., & Pareja-Galeano, H. (2021). Vitamin D and cardiovascular health. *Clinical Nutrition*, *40*(5), 2946-2957.

Dawson-Hughes, B. (2017). Vitamin D and muscle function. *The Journal of steroid biochemistry and molecular biology*, *173*, 313-316.

Buttriss, J. L., & Lanham-New, S. A. (2020). Is a vitamin D fortification strategy needed?. *Nutrition bulletin*, *45*(2), 115.

Ratajczak, A. E., Zawada, A., Rychter, A. M., Dobrowolska, A., & Krela-Kaźmierczak, I. (2021). Milk and dairy products: good or bad for human bone? practical dietary recommendations for the prevention and management of osteoporosis. *Nutrients*, *13*(4), 1329.

Cashman, K. D. (2024). Vitamin D fortification of foods–sensory, acceptability, cost, and public acceptance considerations. *The Journal of Steroid Biochemistry and Molecular Biology*, *239*, 106494.

Pellegrino, L., Marangoni, F., Muscogiuri, G., D'Incecco, P., Duval, G. T., Annweiler, C., & Colao, A. (2021). Vitamin D fortification of consumption cow's milk: Health, nutritional and technological aspects. A multidisciplinary lecture of the recent scientific evidence. *Molecules*, *26*(17), 5289.

Pellegrino, L., Marangoni, F., Muscogiuri, G., D'Incecco, P., Duval, G. T., Annweiler, C., & Colao, A. (2021). Vitamin D fortification of consumption cow's milk: Health, nutritional and technological aspects. A multidisciplinary lecture of the recent scientific evidence. *Molecules*, *26*(17), 5289.

Amrein, K., Scherkl, M., Hoffmann, M., Neuwersch-Sommeregger, S., Köstenberger, M., Tmava Berisha, A., ... & Malle, O. (2020). Vitamin D deficiency 2.0: an update on the current status worldwide. *European journal of clinical nutrition*, *74*(11), 1498-1513.

Jääskeläinen, T., Itkonen, S. T., Lundqvist, A., Erkkola, M., Koskela, T., Lakkala, K., ... & Lamberg-Allardt, C. (2017). The positive impact of general vitamin D food fortification policy on vitamin D status in a representative adult Finnish population: evidence from an 11-y follow-up based on standardized 25-hydroxyvitamin D data. *The American journal of clinical nutrition*, *105*(6), 1512-1520.

Itkonen, S. T., Andersen, R., Björk, A. K., Brugård Konde, Å., Eneroth, H., Erkkola, M., ... & Lamberg-Allardt, C. J. (2021). Vitamin D status and current policies to achieve adequate vitamin D intake in the Nordic countries. *Scandinavian journal of public health*, *49*(6), 616-627.

Vatanparast, H., & Longworth, Z. L. (2023). How does Canada's new vitamin D fortification policy affect the high prevalence of inadequate intake of the vitamin?. *Applied Physiology, Nutrition, and Metabolism*, *48*(11), 870-875.

Siddique, M. T., Hafeez, A., Mearaj, S., & Abbas, S. (2021). Overview of Vitamin D, its status and consequences: Challenges and prospects for Pakistani population: A Review. *Biomedical Letters*, *7*(1), 1-1.

Niedermaier, T., Gredner, T., Kuznia, S., Schöttker, B., Mons, U., Lakerveld, J., ... & PEN-Consortium. (2022). Vitamin D food fortification in European countries: the underused potential to prevent cancer deaths. *European Journal of Epidemiology*, *37*(4), 309-320.

Lavelli, V., D'Incecco, P., & Pellegrino, L. (2021). Vitamin D incorporation in foods: Formulation strategies, stability, and bioaccessibility as affected by the food matrix. *Foods*, *10*(9), 1989.

Pellegrino, L., Marangoni, F., Muscogiuri, G., D'Incecco, P., Duval, G. T., Annweiler, C., & Colao, A. (2021). Vitamin D fortification of consumption cow's milk: Health, nutritional and technological aspects. A multidisciplinary lecture of the recent scientific evidence. *Molecules*, *26*(17), 5289.

McCourt, A. F., & O'Sullivan, A. M. (2022). Using food fortification to improve vitamin D bioaccessibility and intakes. *Proceedings of the Nutrition Society*, *81*(1), 99-107.

Gallo, S., McDermid, J. M., Al-Nimr, R. I., Hakeem, R., Moreschi, J. M., Pari-Keener, M., ... & Cheng, F. W. (2020). Vitamin D supplementation during pregnancy: an evidence analysis center systematic review and meta-analysis. *Journal of the Academy of Nutrition and Dietetics*, *120*(5), 898-924.

Dunlop, E., Kiely, M., James, A. P., Singh, T., & Black, L. J. (2020). Efficacy of vitamin D food fortification and biofortification in children and adults: a systematic review protocol. *JBI evidence synthesis*, *18*(12), 2694-2703.

Augustine, L. F., Nair, K. M., & Kulkarni, B. (2021). Sun exposure as a strategy for acquiring vitamin D in developing countries of tropical region: Challenges & way forward. *Indian Journal of Medical Research*, *154*(3), 423-432.

Wimalawansa, S. J. (2023). Physiological basis for using vitamin D to improve health. *Biomedicines*, *11*(6), 1542.

Zhou, H., Zheng, B., Zhang, Z., Zhang, R., He, L., & McClements, D. J. (2021). Fortification of plant-based milk with calcium may reduce vitamin D bioaccessibility: An in vitro digestion study. *Journal of Agricultural and Food Chemistry*, *69*(14), 4223-4233.

Cashman, K. D. (2022). 100 YEARS OF VITAMIN D: Global differences in vitamin D status and dietary intake: a review of the data. *Endocrine Connections*, *11*(1).

Maurya, V. K., Bashir, K., & Aggarwal, M. (2020). Vitamin D microencapsulation and fortification: Trends and technologies. *The Journal of steroid biochemistry and molecular biology*, *196*, 105489.

El Hoss, K., Salla, M., Khaled, S., Krayem, M., Hassan, H., & El Khatib, S. (2023). Update on vitamin D deficiency and its impact on human health major challenges & technical approaches of food fortification. *Journal of Agriculture and Food Research*, *12*, 100616.

El Hoss, K., Salla, M., Khaled, S., Krayem, M., Hassan, H., & El Khatib, S. (2023). Update on vitamin D deficiency and its impact on human health major challenges & technical approaches of food fortification. *Journal of Agriculture and Food Research*, *12*, 100616.

Benedik, E. (2021). Sources of vitamin D for humans. *International Journal for Vitamin and Nutrition Research*.

Moyersoen, I., Devleesschauwer, B., Dekkers, A., De Ridder, K., Tafforeau, J., Van Camp, J., ... & Lachat, C. (2017). Intake of fat-soluble vitamins in the Belgian population: adequacy and contribution of foods, fortified foods and supplements. *Nutrients*, *9*(8), 860.

Jaaskelainen, T., Itkonen, S. T., Lundqvist, A., Erkkola, M., Koskela, T., Lakkala, K., ... & Harkanen, T. (2017). hman KD, Mannisto S and Lamberg-Allardt C, The positive impact of general vitamin D food fortification policy on vitamin D status in a representative adult Finnish population: evidence from an 11-y follow-up based on standardized 25-hydroxyvitamin D data. *Am. J. Clin. Nutr*, *105*, 1512-1520.

Black, L. J., Walton, J., Flynn, A., Cashman, K. D., & Kiely, M. (2015). Small increments in vitamin D intake by Irish adults over a decade show that strategic initiatives to fortify the food supply are needed. *The Journal of nutrition*, *145*(5), 969-976

Rusińska, A., Płudowski, P., Walczak, M., Borszewska-Kornacka, M. K., Bossowski, A., Chlebna-Sokół, D., ... & Zygmunt, A. (2018). Vitamin D supplementation guidelines for general population and groups at risk of vitamin D deficiency in Poland—recommendations of the polish Society of Pediatric Endocrinology and Diabetes and the expert panel with participation of national specialist consultants and

representatives of scientific societies—2018 update. *Frontiers in endocrinology*, *9*, 246.

Spiro, A., & Buttriss, J. (2014). Vitamin D: an overview of vitamin D status and intake in E urope. *Nutrition bulletin*, *39*(4), 322-350.

Maurya, V. K., Bashir, K., & Aggarwal, M. (2020). Vitamin D microencapsulation and fortification: Trends and technologies. *The Journal of steroid biochemistry and molecular biology*, *196*, 105489.

Itkonen, S. T., Andersen, R., Björk, A. K., Brugård Konde, Å., Eneroth, H., Erkkola, M., ... & Lamberg-Allardt, C. J. (2021). Vitamin D status and current policies to achieve adequate vitamin D intake in the Nordic countries. *Scandinavian journal of public health*, *49*(6), 616-627.

Grégoire-Pelchat, P., Alos, N., Ribault, V., Pastore, Y., Robitaille, N., & Mailhot, G. (2018). Vitamin D intake and status of children with sickle cell disease in Montreal, Canada. *Journal of pediatric hematology/oncology*, *40*(8), e531-e536.

Lamberg-Allardt, C. (2006). Vitamin D in foods and as supplements. Progress in biophysics and molecular biology, 92(1), 33-38.

Biasucci, G., Donini, V., & Cannalire, G. (2024). Rickets Types and Treatment with Vitamin D and Analogues. *Nutrients*, *16*(3), 416.

Souza, S. V., Borges, N., & Vieira, E. F. (2022). Vitamin d-fortified bread: Systematic review of fortification approaches and clinical studies. *Food chemistry*, *372*, 131325.

Niedermaier, T., Gredner, T., Kuznia, S., Schöttker, B., Mons, U., Lakerveld, J., ... & PEN-Consortium. (2022). Vitamin D food fortification in European countries: the underused potential to prevent cancer deaths. *European Journal of Epidemiology*, *37*(4), 309-320.

Neill, H. R., Gill, C. I., McDonald, E. J., McRoberts, W. C., & Pourshahidi, L. K. (2021). Vitamin D biofortification of pork may offer a food-based strategy to increase vitamin D intakes in the UK population. *Frontiers in nutrition*, *8*, 777364.

Dehghan, M., Mente, A., Rangarajan, S., Sheridan, P., Mohan, V., Iqbal, R., ... & Yusuf, S. (2018). Association of dairy intake with cardiovascular disease and mortality in 21 countries from five continents (PURE): a prospective cohort study. *The Lancet*, *392*(10161), 2288-2297.

IEER
Excellence in Education & Research