Policy Research Journal
ISSN (E): 3006-7030 ISSN (P) : 3006-7022 Volume 4, Issue 3, 2026

SOIL HEALTH MANAGEMENT AND SUSTAINABLE AGRICULTURAL PRACTICES FOR IMPROVING CROP PRODUCTIVITY AND ECOSYSTEM SERVICES

Zainab Bibi[*1], Shadman Khan[2], Muhammad ikram[3], Muhammad ikhtisham[4], Aiman Akbar[5], Khadija Tul Kubra[6], Noman Basheer[7]

[*1] *Institute of Soil and Environmental Sciences, PMAS Arid agriculture university Rawalpindi*
[2]*Department: Soil and Environmental Sciences, The University of Agriculture Peshawar*
[3]*Department of Soil and environmental science,The University of agriculture peshawar*
[4]*Department soil and environmental sciences, University of Agriculture Peshawar*
[5]*Department of Agronomy, University of Agriculture Faisalabad*
[6]*Department of Agronomy, University of Agriculture Faisalabad Pakistan*
[7]*Institute of Soil and Environmental Sciences, University of Agriculture, Faisalabad, Pakistan.*

[*1]zainabbibizb98@gmail.com, [2]Shadmankhan@aup.edu.pk, [3]m.ikram@aup.edu.pk, [4]muhammadikhtisham2@gmail.com, [5]Aimanakbar043@gmail.com, [6]khadijakhan512@gmail.com, [7]noumanbaloch266@gmail.com

DOI: https://doi.org/10.5281/zenodo.23209234

Keywords
soil health indicators, conservation agriculture, biochar, cover crops, precision agriculture, ecosystem services, soil organic carbon, sustainable soil management.

Article History
Received: 07 January 2026
Accepted: 17 March 2026
Published: 31 March 2026

Copyright @Author
Corresponding Author: *
Zainab Bibi

Abstract
Soil health is fundamental to global ecosystem stability, agricultural sustainability, and food security. This review synthesizes current knowledge on soil health indicators, standardized assessment frameworks, and sustainable agronomic management practices for improving crop productivity and ecosystem service delivery. The scientific paradigm has evolved from the narrow concept of "soil quality" to the holistic, dynamic concept of "soil health," recognizing soil as a self-regulating biological matrix. Key indicators span physical (bulk density, aggregate stability, water infiltration), chemical (pH, cation exchange capacity, available phosphorus), and biological (microbial biomass carbon, mineralizable nitrogen, β-glucosidase activity, metabolic quotient) domains, with soil organic carbon acting as the central cross-cutting variable. Minimum Data Sets within standardized frameworks including CASH, SMAF, SHAPE, HAST, and VESS convert heterogeneous indicator values into actionable management insights while distinguishing inherent from dynamic soil properties. Meta-analytic synthesis reveals that no-till in isolation incurs a 5.7–12% global yield penalty; however, integrated Conservation Agriculture combining zero tillage, residue retention, and crop diversification eliminates this penalty and increases yields in rainfed dryland environments. Leguminous cover crops increase the mineralizable nitrogen pool by 211%, while diversified rotations increase it by 44%. Biochar amendments increase soil organic carbon stocks by 29% and reduce methane emissions by 51.31%. Precision agriculture integrating IoT sensors, AI, and variable-rate technology enables targeted, spatially explicit soil management. Collectively, these

practices enhance provisioning, regulating, and supporting ecosystem services, building agricultural resilience against climate shocks.

INTRODUCTION

Soil health is fundamental to global ecosystem stability, agricultural sustainability, and food security. Historically defined primarily through the narrow lens of crop yield capacity and chemical fertility, the scientific paradigm has undergone a major evolution from "soil quality" which emphasizes functional land-use capabilities to the dynamic, holistic concept of "soil health" (Rehman et al., 2024). Contemporary consensus defines soil health as the continued capacity of soil to function as a vital living ecosystem that sustains biological productivity, maintains air and water quality, and promotes plant, animal, and human health. This conceptual reframing recognizes that soil is not an inert physical medium, but rather a complex, self-regulating biological matrix governed by continuous interactions among mineral phases, organic matter, and subterranean biological communities (Sharma et al., 2024).

The imperative for sustainable soil health management is driven by widespread land degradation, soil erosion, organic matter depletion, and climate variability. Globally, hundreds of millions of hectares of arable land suffer from physical compaction, chemical salinization, nutrient mining, and severe loss of biological diversity (Xing et al., 2025). In regions undergoing intensive agricultural exploitation, excessive tillage and agrochemical over-reliance have decoupled natural biogeochemical cycles, degrading the long-term productive capacity of terrestrial agroecosystems. Addressing these complex challenges requires a structured, tripartite scientific approach encompassing conceptualization, assessment, and management (Alananbeh, 2020). Conceptualization establishes soil as a dynamic biological system; assessment involves measuring sensitive physical, chemical, and biological indicators; and management deploys targeted agronomic interventions to restore functionality (Powlson et al., 2011).

This review synthesizes current knowledge regarding soil health indicators, standardized assessment frameworks, and sustainable agronomic management practices. It evaluates the quantitative impacts of conservation tillage, cover cropping, crop rotations, biochar amendments, and precision digital soil mapping on crop productivity and ecosystem service delivery (Teague & Kreuter, 2020).

2. Key Soil Health Indicators: Physical, Chemical, and Biological Metrics

Assessing soil functionality requires measuring quantifiable indicators sensitive to management changes and environmental disturbances. Soil properties are broadly categorized into physical, chemical, and biological parameters. Among these, soil organic carbon (SOC) acts as the central cross-cutting variable that directly modulates physical structure, chemical buffer capacity, and biological activity (Cardoso et al., 2013).




Figure 1. Conceptual Tripartite Triad of Soil Health: Physical, Chemical, and Biological Indicator Metrics Mediating Agricultural Practices and Global Food Security

2.1 Physical Soil Health Metrics

Soil physical properties govern the structural framework, mechanical strength, aeration, thermal capacity, and hydrologic behavior of the soil matrix. Bulk density serves as a fundamental metric for evaluating soil compaction and macro-porosity. Elevated bulk density values restrict root elongation, reduce pore continuity, and impede gas exchange, creating hypoxic conditions that restrict root respiration. Intensive tillage degrades structural pore networks, whereas conservation tillage and residue retention lower bulk density over time by promoting biological macropore formation (Fine et al., 2017).

Aggregate stability measures the capacity of soil structural units to resist breakdown when exposed to erosive forces such as raindrop impact or water slaking. Structural stabilization relies on micro-aggregates being bound into durable macro-aggregates by organic binding agents, plant root mucilage, fungal hyphae, and biogenic glycoproteins like glomalin, which is secreted by arbuscular mycorrhizal fungi. High aggregate stability improves water infiltration rates and

expands soil water-holding capacity, allowing soils to store plant-available water and mitigate crop drought stress (Norris et al., 2020).

2.2 Chemical Soil Health Metrics

Soil chemical properties regulate the thermodynamic availability, mobility, and exchangeability of essential plant nutrients and potential phytotoxins. Soil pH is widely recognized as the master soil variable due to its profound influence on biogeochemical equilibria, chemical speciation, and biological activity. Deviations from the optimal pH range for most field crops (6.0–7.5) can induce aluminum or manganese toxicities in acidic soils, precipitate insoluble calcium phosphate complexes in alkaline soils, and alter microbial enzymatic reaction kinetics (Willoughby et al., 2023).

Cation exchange capacity (CEC) measures the density of negative surface charges on clay minerals and organic matter complexes, quantifying the soil's ability to retain exchangeable cations (Ca^{2+}, Mg^{2+}, K^{+}, Na^{+}) against hydrological leaching. Concurrently, electrical conductivity (EC) monitors total soluble salt concentrations; elevated EC indicates soil salinization, which exerts osmotic stress on crop roots and degrades soil structural integrity (Raghavendra et al., 2020). Nutrient availability metrics including total organic carbon, total nitrogen, Olsen phosphorus, and exchangeable potassium provide quantitative measures of chemical fertility and long-term nutrient-supplying capacity (Fine et al., 2017).

2.3 Biological and Biochemical Indicators

Soil biological indicators offer high sensitivity and early-warning capabilities, reflecting shifts in land management long before alterations in total soil organic carbon stocks become analytically detectable. Microbial biomass carbon (MBC) represents the active living fraction of soil organic matter, serving as both a sensitive sink and source of plant-available nutrients (García & Hernández, 1997). Potentiated mineralizable nitrogen (PMN) measures the labile organic nitrogen pool capable of microbial conversion into plant-available inorganic forms (NH_4^+ and NO_3^-), reflecting active biological nitrogen cycling (Zaghloul et al., 2020). Respiration rates and metabolic quotients describe the metabolic state and stress level of soil microbial communities. The metabolic quotient (qCO_2), defined as the ratio of basal microbial respiration to total microbial biomass carbon, reflects microbial carbon-use efficiency; elevated qCO_2 values indicate microbial metabolic stress or ecosystem disturbance. Conversely, the microbial quotient (qMIC), defined as the ratio of microbial carbon to total organic carbon, reflects substrate availability and the efficiency with which organic inputs are incorporated into microbial biomass (Dahlhoff, 2004).

Extracellular soil enzymes synthesized by microorganisms and plant roots catalyze the rate-limiting steps of organic matter decomposition and nutrient mineralization. The enzyme β-glucosidase catalyzes the hydrolysis of terminal cellobiose residues into labile glucose, serving as a primary bio-indicator of labile carbon cycling (Bending et al., 2004). Ratios of β-glucosidase activity to soil organic carbon exceeding 17 signal net carbon sequestration trends, whereas ratios below 17 indicate net organic carbon mineralization. Similarly, N-acetyl-β-glucosaminidase (NAG) cleaves fungal chitin to release organic nitrogen, while phosphomonoesterase and arylsulfatase hydrolyze ester bonds to release orthophosphate and sulfate, controlling biological phosphorus and sulfur availability (Vittori Antisari et al., 2021). Larger soil organisms, including earthworms, collembolans, and beneficial nematodes, physically process organic residues, create macro-burrows that enhance aeration, and modulate subterranean food webs (Damásio et al., 2007).

Table 1: Diagnostic Classification and Functional Roles of Key Soil Health Indicators

Indicator Domain	Parameter	Primary Measurement Method	Functional Role & Sensitivity
Physical	Bulk Density (ρ_b)	Dry core mass per unit volume (g cm^{-3})	Evaluates compaction, mechanical root impedance, and macro-porosity.
Physical	Aggregate Stability	Wet-sieving / Mean weight diameter (MWD)	Measures structural resistance to water erosion, slaking, and surface crusting.
Physical	Water Infiltration Rate	Double-ring infiltrometer / Saturated hydraulic conductivity	Quantifies surface water entry capacity, runoff risk, and topsoil erosion potential.
Chemical	Soil pH	Potentiometric determination (1: 1 H_2O or $CaCl_2$)	Regulates nutrient chemical speciation, trace element toxicity, and microbial kinetics.
Chemical	Cation Exchange Capacity	Ammonium acetate displacement ($cmol_c$ kg^{-1})	Quantifies charge density available to retain plant-essential exchangeable cations.
Chemical	Available Phosphorus	Olsen-P / Bray-P bicarbonate extraction	Measures plant-available orthophosphate concentrations in soil solution.
Biological	Microbial Biomass C	Chloroform fumigation-extraction / PLFA	Quantifies active living microbial carbon pool and labile nutrient reservoir.
Biological	Mineralizable Nitrogen	Anaerobic incubation / Citrate extractable protein	Measures biological capacity to mineralize labile organic N into crop-available forms.
Biological	β-Glucosidase (BG)	Colorimetric assay (p-nitrophenyl-β-D-glucoside)	Early-warning bio-indicator of labile carbon turnover and decomposition kinetics.
Biological	Metabolic Quotient (qCO_2)	Ratio of basal respiration to total microbial biomass C	Measures microbial carbon utilization efficiency and metabolic stress response.

3. Soil Health Assessment Frameworks and Minimum Data Sets

Due to the labor intensity and analytical expense of comprehensive testing, evaluating soil health across agricultural landscapes requires selecting targeted Minimum Data Sets (MDS) within standardized scoring frameworks. An MDS comprises a reduced, non-redundant set of physical, chemical, and biological parameters selected through statistical dimensionality reduction such as principal component analysis or factor analysis to capture essential soil functions (Andrews et al., 2024). Standard agricultural Minimum Data Sets typically integrate core variables including soil pH, total organic carbon, total nitrogen, available phosphorus, mineralizable nitrogen, bulk density, aggregate stability, and microbial respiration (Doran & Parkin, 1997).

Once MDS indicator values are measured, standard assessment frameworks convert raw heterogeneous units into non-linear, unitless scores ranging from 0 to 100 based on established ecological baseline curves. Scoring curves are categorized into three primary functional relationships: "more is better" (e.g., organic matter,

aggregate stability), "less is better" (e.g., bulk density, electrical conductivity), and "optimum range" (e.g., soil pH, available phosphorus) (Hughes et al., 2023).

Several standardized index tools convert indicator scores into actionable soil management insights:

- The Comprehensive Assessment of Soil Health (CASH), developed by Cornell University, integrates physical, chemical, and biological metrics into localized index scores, pairing diagnostic sub-scores with specific management strategies (Yuan et al., 2026).
- The Soil Management Assessment Framework (SMAF) uses algorithms calibrated for soil climate regimes, mineralogy, slope, and crop requirements to evaluate management effects on specific soil functions.
- The Soil Health Assessment Protocol and Evaluation (SHAPE) incorporates spatial peer-group reference datasets across pedoclimatic gradients, ensuring soil health scores account for inherent pedogenic factors (Nunes et al., 2021).
- The Haney Soil Test (HAST) evaluates dual weak-acid extractants paired with 24-hour microbial respiration burst assays to quantify labile organic carbon and nitrogen pools for dynamic fertilizer calibrations.
- The Visual Evaluation of Soil Structure (VESS) provides a rapid field methodology that visually rates aggregate size, macroporosity, and root penetration without laboratory equipment (Das et al., 2026).

Accurately interpreting soil health frameworks requires distinguishing between *inherent* soil properties (genoforms) and *dynamic* soil properties (phenoforms). Inherent properties are fixed by pedogenic factors such as parent material, texture, climate, topography, and time; for example, a sandy Entisol cannot match the native CEC or water-holding capacity of a clay-rich Vertisol (Mponela et al., 2026). Dynamic properties are responsive to human management decisions over shorter timescales. Soil health assessments must evaluate dynamic indicator trajectories relative to localized pedoclimatic reference states rather than attempting to apply universal absolute targets across diverse soil types (Jian et al., 2020).

4. Sustainable Agricultural Practices and Agronomic Productivity

Reversing soil degradation requires transitioning from intensive conventional management to conservation-oriented systems that restore soil functionality while sustaining crop yields (Lal, 2008).

4.1 Conservation Agriculture: The Tripartite Model

Conservation Agriculture (CA) is built on three linked agronomic principles: minimal mechanical soil disturbance (no-till or reduced tillage), permanent organic soil cover via crop residue retention or cover crops, and diversified crop rotations involving three or more species (Zhang et al., 2024). Conventional moldboard plowing shearing breaks soil aggregates, exposes protected organic matter to rapid microbial oxidation, accelerates atmospheric carbon loss, and destroys subterranean fungal networks. In contrast, continuous no-till preserves soil structural integrity, protects fungal hyphae, reduces surface bulk density over time, and concentrates organic matter in the upper soil profile (Campobasso et al., 2025).




Figure 2. Consolidated Framework of Soil Health Management Principles: Interlinking Minimal Soil Disturbance, Permanent Surface Cover, Crop Diversity, Year-Round Roots, and Livestock Integration

4.2 Meta-Analytic Synthesis of No-Till Dynamics and Yield Trade-Offs

While no-till improves physical and biological soil health metrics, long-term global meta-analyses highlight spatial and agronomic nuances in crop yield responses. Synthesizing over 600 peer-reviewed studies comprising more than 5,000 paired yield comparisons demonstrates that implementing no-till *in isolation* results in an average global yield penalty of 5.7% to 12% compared to conventional tillage, particularly when synthetic nitrogen inputs are limited (Adil, 2026).

This yield deficit under standalone no-till stems from specific physical and biological mechanisms: early-phase surface compaction, lower soil temperatures under surface residue in cool humid climates, delayed seed germination, early weed competition, and nitrogen immobilization caused by high-C/N surface crop residues (Cui et al., 2024). Cereal crops display notable yield reductions under standalone no-till, with average yield declines of 7.6% for maize, 7.5% for rice, and 2.6% for wheat. Conversely, oilseed, cotton, and leguminous crop yields show minimal impact, matching conventional tillage productivity under

standalone no-till management (Von Cossel et al., 2025).

Crucially, global synthesis reveals that the yield penalty associated with no-till is minimized or reversed when all three Conservation Agriculture principles zero tillage, permanent crop residue retention, and crop rotation are implemented simultaneously. In rainfed, arid and semi-arid agroecosystems, fully integrated Conservation Agriculture consistently outperforms conventional tillage, increasing crop yields (Lin et al., 2023). Surface residue retention acts as protective mulch, suppressing evaporation, moderating soil temperatures, enhancing infiltration rates, and conserving critical root-zone water stores during mid-season dry spells. Furthermore, initial yield declines under no-till dissipate over time; crop yields under conservation management typically equal or exceed conventional tillage benchmarks after 3 to 10 years of continuous practice as soil aggregate networks, macroporosity, and natural nutrient cycles recover (Song et al., 2026).

4.3 Cover Crops and Crop Diversity Effects

Integrating cover crops into fallow windows maintains continuous living root systems that secrete diverse root exudates, sustaining active rhizosphere microbial communities throughout the year. Leguminous cover crops establish symbiotic relationships with N_2-fixing rhizobia bacteria, adding biologically fixed nitrogen to the soil system while providing low-C/N organic residues that mineralize rapidly (Beillouin et al., 2021).

Synthesizing multi-site experimental data shows that incorporating leguminous cover crops increases the mineralizable nitrogen (PMN) pool by 211% compared to fallow controls. Diversified crop rotations containing three or more distinct species increase the PMN pool by 44% compared to continuous monocultures, whereas no-till management alone increases PMN by 13%. Diverse crop rotations disrupt host-specific pathogen cycles, lower weed pressure, introduce varied root structures, and promote microbial functional redundancy, which stabilizes long-term crop yields during climatically stressful years (Quintarelli et al., 2022).

Table 2: Agronomic Impact and Soil Health Responses to Conservation Management Practices

Management Practice	Primary Soil Impact Mechanism	Quantitative Soil Metric Response	Global Yield Impact & Context
No-Till Alone	Minimizes mechanical shear; protects aggregate structures and hyphae.	$+13\%$ Potentiated Mineralizable N; topsoil organic C accumulation.	-5.7% to -12% average yield loss; yield penalties highest in humid regions.
Integrated Conservation Agriculture	Combines zero-tillage, surface residue mulch, and crop rotation.	Maximizes macro-aggregate stability, macroporosity, and infiltration.	Eliminates yield penalties; increases crop yields in rainfed dryland environments.
Leguminous Cover Crops	Biological N_2 fixation; supplies low-C/N labile organic matter to soil.	$+211\%$ increase in Potentiated Mineralizable Nitrogen pool.	Increases subsequent main crop yields; offsets synthetic N fertilizer demand.
Diversified Rotations ($\geq$ 3 crops)	Expands root exudate diversity; breaks host-pathogen cycles.	$+44\%$ increase in Potentiated Mineralizable Nitrogen pool.	Improves multi-year yield stability; reduces temporal yield variance.

5. Soil Organic Amendments and Biochar Dynamics

Applying exogenous organic amendments such as composts, animal manures, and biochar serves as an effective strategy for accelerating the recovery of degraded agricultural soils (Gross et al., 2024).

5.1 Biochar Pyrolysis, Stability, and SOC Sequestration

Biochar is a carbon-rich, highly porous, recalcitrant material produced by pyrolyzing biomass feedstocks including crop residues, forestry residues, and livestock manures at temperatures ranging from 350°C to 700°C under limited oxygen supply. Thermochemical conversion converts labile aliphatic carbon into highly condensed aromatic ring structures that resist microbial mineralization, creating a long-term terrestrial carbon sink (Li & Tasnady, 2023). Global meta-analyses of long-term field studies show that applying biochar at rates between 1 and 100 Mg ha^{-1} increases total soil organic carbon stocks by an average of 29%, corresponding to a mean carbon accumulation of 13.0 Mg SOC ha^{-1}. In controlled incubation and pot studies, biochar additions raise soil organic carbon content by 75%, corresponding to a mean increase of 6.3 g SOC kg^{-1} soil (Luo et al., 2023). Wood- and plant-derived biochars produced at higher pyrolysis temperatures display greater recalcitrance and carbon sequestration capacity than manure-derived biochars due to their higher C/N ratios and aromaticity. The magnitude of organic carbon retention is also governed by soil texture; medium- to fine-textured clay soils stabilize biochar carbon more effectively than coarse sandy soils through organo-mineral interactions (Rodrigues et al., 2023).

5.2 Synergistic Effects of Biochar-Compost Co-Application

Applying raw, unamended biochar directly to soil can occasionally cause short-term crop nitrogen stress due to the sorption of plant-available mineral nitrogen onto fresh biochar surfaces. Co-application of biochar with high-nutrient organic fertilizers, or processing biochar through co-composting mixtures, mitigates this limitation and generates functional synergies (Pamuru et al., 2026).

During co-composting, porous biochar serves as a structural bulking agent that increases aeration, reduces anaerobic micro-zones, and accelerates organic matter stabilization. Meta-analysis of biochar-amended composting processes demonstrates significant improvements in compost quality: the germination index increases by 25.57% (indicating reduced phytotoxicity), nitrate-nitrogen ($NO_3^- - N$) concentrations increase by 32.54%, ammonium-nitrogen ($NH_4^+ - N$) volatilization losses decrease by 34.96%, and the final C/N ratio drops by 5.44% (Wang et al., 2025).

When applied to field soils, biochar-compost mixtures deliver performance superior to sole biochar or sole compost applications. The internal pore network of biochar absorbs labile humic compounds, amino acids, and inorganic nutrients from the compost matrix, forming organo-mineral coatings on biochar surfaces. This process reduces nutrient leaching, increases cation exchange capacity, enhances moisture retention, and supports plant growth, particularly in highly weathered tropical soils prone to rapid organic matter decomposition (Luo et al., 2026).

5.3 Greenhouse Gas Mitigation and Nutrient Retention Mechanisms

Biochar amendments alter nitrogen and carbon biogeochemistry, reducing gaseous emissions from agroecosystems:

- **Methane (CH_4) Suppression**: Incorporating biochar into composting systems reduces overall methane emissions by 51.31%, with reductions exceeding 86% when biochar inclusion rates exceed 12% by weight. Biochar improves compost structural aeration, suppressing methanogenic archaea while encouraging methanotrophic bacterial activity that oxidizes produced methane into carbon dioxide (Rubin et al., 2020).
- **Nitrous Oxide (N_2O) Mitigation**: In saturated agricultural soils, biochar reduces nitrous oxide emissions by acting as an electron shuttle during microbial denitrification, facilitating the complete enzymatic reduction of N_2O to inert N_2 gas.
- **Ammonia (NH_3) Volatilization Reduction**: The functional groups and high cation exchange capacity of lower-temperature biochars absorb ammonium ions (NH_4^+), limiting free ammonia

volatilization from applied animal manures and slurry (Sun et al., 2022).

6. Precision Agriculture, Digital Soil Mapping, and Sensor Technologies

Modern soil management increasingly relies on spatial sensing tools to address natural soil variability across field landscapes. Precision agriculture combines high-resolution data acquisition with targeted input management to optimize resource use efficiency (Paramesh et al., 2023).

6.1 Proximal and Remote Sensing for High-Resolution Soil Character characterization

Digital Soil Mapping (DSM) relates observed soil properties with spatial environmental covariates such as digital elevation models, climatic rasters, and spectral remote sensing imagery to predict soil metrics across landscape scales (Ciampalini et al., 2015). Proximal soil sensing technologies, such as field-portable visible and near-infrared (VIS-NIR) and mid-infrared (MIR) diffuse reflectance spectrometers, enable rapid, non-destructive estimation of soil organic carbon, moisture, total nitrogen, and clay mineralogy. Field-deployed electromagnetic induction (EMI) sensors measure apparent electrical conductivity (EC_a), mapping spatial variations in soil salinity, texture, and moisture saturation (Duda et al., 2017).

Complementing ground-based sensors, remote sensing platforms including satellite constellations (e.g., Sentinel-2, Landsat-9) and unmanned aerial vehicles (UAVs) equipped with multispectral or hyperspectral payloads evaluate surface soil organic carbon patterns, bare-soil spectral signatures, and vegetation indexes (e.g., NDVI) to monitor crop canopy health and stress (Koumoulidis et al., 2024).

6.2 AI, Machine Learning, and IoT Sensor Meshes in Variable-Rate Management

Integrating Internet of Things (IoT) sensor arrays with artificial intelligence (AI) and machine learning (ML) algorithms enables continuous real-time monitoring of soil micro-climates and nutrient dynamics. In-situ IoT sensor nodes deployed across fields transmit real-time telemetry detailing soil moisture, matric potential, soil temperature, pH, and nitrate concentrations to cloud processing platforms (Yaqoob et al., 2026).

Machine learning models such as Random Forest, Support Vector Machines, and Deep Neural Networks process spatial covariate layers, remote sensing imagery, and real-time IoT data to generate high-resolution field prescription maps (Gulve, 2023). Variable-Rate Technology (VRT) equipment uses these prescription maps to continuously modulate application rates of lime, nitrogen, phosphorus, and irrigation water across heterogeneous fields. This targeted spatial management corrects localized nutrient deficiencies, avoids over-fertilization, reduces chemical leaching into aquifers, and lowers input costs (Nicola et al., 2026).

6.3 Bridging Spatial Scale Gaps between Digital Soil Maps and Field-Level Realities

A persistent challenge in precision agriculture is resolving spatial scale mismatches between regional digital soil maps and field-level operational realities. Coarse regional digital maps (grid resolutions of 30 m to 250 m) frequently fail to capture sub-meter micro-topographic variations, localized compaction zones, or field-scale drainage tile effects (Vergopolan et al., 2020).

Addressing this spatial gap requires multi-scale data fusion techniques that combine broad satellite imagery with high-density proximal sensor sweeps and targeted core sampling. Establishing standardized calibration protocols such as those evaluated by the North American Project to Evaluate Soil Health Measurements (NAPESHM) ensures that rapid spectral assays and smart sensor applications produce reliable soil health data that align with standard laboratory reference methods (Chen et al., 2026).

7. Ecosystem Services and Environmental Resilience

Sustaining healthy agricultural soils delivers vital ecosystem services that extend beyond crop biomass production, contributing to broad

environmental sustainability. Ecosystem services provided by managed soils fall into three main functional domains:

1. **Provisioning Services**: Healthy soils maintain agricultural productivity by providing structural root support, regulating essential plant nutrient supply, and expanding available water storage. These functions stabilize multi-year crop yields and improve food security under variable climatic conditions (Biggs et al., 2012).

2. **Regulating Services**: Managed soils contribute to climate change mitigation by sequestering carbon into mineral-associated organic matter and stable humic fractions, removing CO_2 from the atmosphere. Improved macro-porosity and structural aggregate stability enhance rain infiltration rates, reducing surface runoff, peak flood volumes, and water-driven soil erosion. Furthermore, active soil matrices serve as natural filters, adsorbing heavy metals, trapping agricultural sediments, and metabolizing pesticide residues before contaminants reach surface waters or groundwater aquifers (Biggs et al., 2015).

3. **Supporting Services**: Soil health management preserves biological diversity within subterranean environments, supporting food webs, maintaining microbial functional redundancy, and driving fundamental biogeochemical cycles (C, N, P, S).

By enhancing these regulating, supporting, and provisioning functions, sustainable soil management increases agricultural resilience against climate shocks, reducing crop yield volatility during extreme weather events (Oliver et al., 2015).

Conclusion

This review demonstrates that soil health management represents a paradigm shift from narrow yield-focused approaches toward holistic, biologically grounded stewardship of agricultural ecosystems. The tripartite framework of conceptualization, assessment, and management provides a structured pathway for addressing widespread land degradation, organic matter depletion, and climate variability. Physical, chemical, and biological indicators integrated through Minimum Data Sets and standardized scoring frameworks such as CASH, SMAF, SHAPE, HAST, and VESS offer sensitive, early-warning diagnostics of management-induced changes, with soil organic carbon serving as the central cross-cutting variable modulating structure, fertility, and biological activity. The evidence synthesized here reveals that no-till alone is insufficient; rather, the full integration of Conservation Agriculture principles zero tillage, permanent residue retention, and diversified rotations eliminates yield penalties and enhances productivity, particularly in rainfed arid and semi-arid agroecosystems. Leguminous cover crops and diversified rotations substantially augment labile nitrogen pools and improve multi-year yield stability. Biochar, especially when co-applied with compost, enhances carbon sequestration, reduces greenhouse gas emissions, and improves nutrient retention. Precision agriculture technologies, including proximal and remote sensing, IoT sensor networks, and AI-driven variable-rate management, bridge spatial scale gaps and enable targeted, resource-efficient interventions. Collectively, these practices enhance provisioning, regulating, and supporting ecosystem services, increasing agricultural resilience against climate shocks while reducing environmental footprints. Future research must prioritize long-term, multi-location field studies, standardized calibration protocols, and farmer-feasible assessment tools to translate scientific insights into actionable, site-specific management recommendations that sustain soil health across diverse pedoclimatic regions.

REFERENCES

Adil, M. (2026). No-Till Legume Cover Crops Enhance Soil Carbon, Mitigate Greenhouse Gas Emissions, and Increase Yield in Dryland Wheat: A Global Meta-Analysis. *Land Degradation & Development*.

Alananbeh, K. (2020). Soil health and sustainable agriculture. *Sustainability*.

Andrews, S. S., Karlen, D. L., & Cambardella, C. A. (2004). The soil management assessment framework: a quantitative soil quality evaluation method. *Soil Science Society of America Journal*, *68*(6), 1945-1962.

Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V., & Makowski, D. (2021). Positive but variable effects of crop diversification on biodiversity and ecosystem services. *Global change biology*, *27*(19), 4697-4710.

Bending, G. D., Turner, M. K., Rayns, F., Marx, M. C., & Wood, M. (2004). Microbial and biochemical soil quality indicators and their potential for differentiating areas under contrasting agricultural management regimes. *Soil Biology and Biochemistry*, *36*(11), 1785-1792.

Biggs, R., Schlüter, M., & Schoon, M. L. (Eds.). (2015). Principles for building resilience: sustaining ecosystem services in social-ecological systems.

Biggs, R., Schlüter, M., Biggs, D., Bohensky, E. L., BurnSilver, S., Cundill, G., ... & West, P. C. (2012). Toward principles for enhancing the resilience of ecosystem services. *Annual review of environment and resources*, *37*, 421-448.

Campobasso, A. A., Frem, M., Petrontino, A., Tricarico, G., & Bozzo, F. (2025). Classification, evaluation and adoption of innovation: A systematic review of the agri-food sector. *Agriculture*, *15*(17), 1845.

Cardoso, E. J. B. N., Vasconcellos, R. L. F., Bini, D., Miyauchi, M. Y. H., Santos, C. A. D., Alves, P. R. L., ... & Nogueira, M. A. (2013). Soil health: looking for suitable indicators. What should be considered to assess the effects of use and management on soil health?. *Scientia agricola*, *70*(4), 274-289.

Chen, J., Zhang, X., Fan, Z., Liu, T., Xiao, W., Sun, Q., ... & Huang, Z. (2026). High-resolution, multi-depth mapping of soil bulk density and pH in China's forests using machine learning. *Earth System Science Data*, *18*(6), 4451-4474.

Ciampalini, A., André, F., Garfagnoli, F., Grandjean, G., Lambot, S., Chiarantini, L., & Moretti, S. (2015). Improved estimation of soil clay content by the fusion of remote hyperspectral and proximal geophysical sensing. *Journal of Applied Geophysics*, *116*, 135-145.

Cui, Y., Zhang, W., Zhang, Y., Liu, X., Zhang, Y., Zheng, X., ... & Zou, J. (2024). Effects of no-till on upland crop yield and soil organic carbon: a global meta-analysis. *Plant and Soil*, *499*(1), 363-377.

Dahlhoff, E. P. (2004). Biochemical indicators of stress and metabolism: applications for marine ecological studies. *Annu. Rev. Physiol.*, *66*(1), 183-207.

Damásio, J. B., Barata, C., Munné, A., Ginebreda, A., Guasch, H., Sabater, S., ... & Porte, C. (2007). Comparing the response of biochemical indicators (biomarkers) and biological indices to diagnose the ecological impact of an oil spillage in a Mediterranean river (NE Catalunya, Spain). *Chemosphere*, *66*(7), 1206-1216.

Das, S., Gamage, L. K. H., Stephenson, M., Acharya, U., Daigh, A. L. M., & Maharjan, B. (2026). A two-tier, reference-informed framework for soil health assessment: Minimum indicator sets and ecologically dynamic thresholds. *Agrosystems, Geosciences & Environment*, *9*(3), e70449.

Doran, J. W., & Parkin, T. B. (1997). Quantitative indicators of soil quality: a minimum data set. *Methods for assessing soil quality*, *49*, 25-37.

Duda, B. M., Weindorf, D. C., Chakraborty, S., Li, B., Man, T., Paulette, L., & Deb, S. (2017). Soil characterization across catenas via advanced proximal sensors. *Geoderma*, *298*, 78-91.

Fine, A. K., Van Es, H. M., & Schindelbeck, R. R. (2017). Statistics, scoring functions, and regional analysis of a comprehensive soil health database. *Soil Science Society of America Journal*, *81*(3), 589-601.

Fine, A. K., Van Es, H. M., & Schindelbeck, R. R. (2017). Statistics, scoring functions, and regional analysis of a comprehensive soil health database. *Soil Science Society of America Journal*, *81*(3), 589-601.

García, C., & Hernández, T. (1997). Biological and biochemical indicators in derelict soils subject to erosion. *Soil Biology and Biochemistry*, *29*(2), 171-177.

Gross, A., Bromm, T., Polifka, S., Fischer, D., & Glaser, B. (2024). Long-term biochar and soil organic carbon stability–Evidence from field experiments in Germany. *Science of the Total Environment*, *954*, 176340.

Gulve, M. (2023). Dr. Ramesh Gawade Dr. Ashish Jadhav Mr. Vaibhav wardule.

Hughes, H. M., Koolen, S., Kuhnert, M., Baggs, E. M., Maund, S., Mullier, G. W., & Hillier, J. (2023). Towards a farmer-feasible soil health assessment that is globally applicable. *Journal of Environmental Management*, *345*, 118582.

Jian, J., Du, X., & Stewart, R. D. (2020). A database for global soil health assessment. *Scientific Data*, *7*(1), 16.

Koumoulidis, D., Efthimiadou, A., Katsenios, N., & Hadjimitsis, D. (2024, September). A Review: Soil properties mapping estimation using remote and proximal sensing data. In *Tenth international conference on remote sensing and geoinformation of the environment (RSCy2024)* (Vol. 13212, pp. 73-86). SPIE.

Lal, R. (2008). Soils and sustainable agriculture. A review. *Agronomy for Sustainable Development*, *28*(1), 57-64.

Li, S., & Tasnady, D. (2023). Biochar for soil carbon sequestration: current knowledge, mechanisms, and future perspectives. *C*, *9*(3), 67.

Lin, B. J., Li, R. C., Liu, K. C., Pelumi Oladele, O., Xu, Z. Y., Lal, R., ... & Zhang, H. L. (2023). Management-induced changes in soil organic carbon and related crop yield dynamics in China's cropland. *Global Change Biology*, *29*(13), 3575-3590.

Luo, C., Zhang, L., Suo, L., & Liang, L. (2026). Synergistic application of biochar and compound microbial inoculant improves the performance of green waste composting. *Journal of Environmental Management*, *397*, 128260.

Luo, L., Wang, J., Lv, J., Liu, Z., Sun, T., Yang, Y., & Zhu, Y. G. (2023). Carbon sequestration strategies in soil using biochar: advances, challenges, and opportunities. *Environmental Science & Technology*, *57*(31), 11357-11372.

Mponela, P., Chimonyo, V. G. P., Chiduwa, M., Kachali, R., Grevulo-Minofu, J., Kawanga, C., ... & Snapp, S. (2026). Soil-health frameworks in agri-food systems. A review: Mponela et al. *Agronomy for Sustainable Development*, *46*(3), 34.

Nicola, F., Giuseppe, T., Spagnuolo, M., Giuseppe, S., Caria, M., & Giampaolo, S. (2026). Innovations in Robots for Weed and Pest Control: A Systematic Review of Cutting-Edge Research. *Machine Learning and Knowledge Extraction*, *8*(2), 51.

Norris, C. E., Bean, G. M., Cappellazzi, S. B., Cope, M., Greub, K. L., Liptzin, D., ... & Honeycutt, C. W. (2020). Introducing the North American project to evaluate soil health measurements. *Agronomy Journal*, *112*(4), 3195-3215.

Nunes, M. R., Veum, K. S., Parker, P. A., Holan, S. H., Karlen, D. L., Amsili, J. P., ... & Moorman, T. B. (2021). The soil health assessment protocol and evaluation applied to soil organic carbon. *Soil Science Society of America Journal*, *85*(4), 1196-1213.

Oliver, T. H., Heard, M. S., Isaac, N. J., Roy, D. B., Procter, D., Eigenbrod, F., ... & Bullock, J. M. (2015). Biodiversity and resilience of ecosystem functions. *Trends in ecology & evolution*, *30*(11), 673-684.

Pamuru, S. T., Brown, B. M., Saltali, O., Farina, A., Davis, A. P., Aydilek, A. H., & Cetin, B. (2026). Effects of compost and biochar co-application on nitrogen leaching and carbon dynamics in soil-water-plant systems. *Journal of Environmental Management*, *398*, 128571.

Paramesh, V., Mohan Kumar, R., Rajanna, G. A., Gowda, S., Nath, A. J., Madival, Y., ... & Toraskar, S. (2023). Integrated nutrient management for improving crop yields, soil properties, and reducing greenhouse gas emissions. *Frontiers in Sustainable Food Systems*, *7*, 1173258.

Powlson, D. S., Gregory, P. J., Whalley, W. R., Quinton, J. N., Hopkins, D. W., Whitmore, A. P., ... & Goulding, K. W. (2011). Soil management in relation to sustainable agriculture and ecosystem services. *Food policy*, *36*, S72-S87.

Quintarelli, V., Radicetti, E., Allevato, E., Stazi, S. R., Haider, G., Abideen, Z., ... & Mancinelli, R. (2022). Cover crops for sustainable cropping systems: a review. *Agriculture*, *12*(12), 2076.

Raghavendra, M., Sharma, M. P., Ramesh, A., Richa, A., Billore, S. D., & Verma, R. K. (2020). Soil health indicators: Methods and applications. In *Soil analysis: recent trends and applications* (pp. 221-253). Singapore: Springer Singapore.

Rehman, A., Farooq, M., Lee, D. J., & Siddique, K. H. (2022). Sustainable agricultural practices for food security and ecosystem services. *Environmental Science and Pollution Research*, *29*(56), 84076-84095.

Rodrigues, L., Budai, A., Elsgaard, L., Hardy, B., Keel, S. G., Mondini, C., ... & Leifeld, J. (2023). The importance of biochar quality and pyrolysis yield for soil carbon sequestration in practice. *European Journal of Soil Science*, *74*(4), e13396.

Rubin, R. L., Anderson, T. R., & Ballantine, K. A. (2020). Biochar simultaneously reduces nutrient leaching and greenhouse gas emissions in restored wetland soils. *Wetlands*, *40*(6), 1981-1991.

Sharma, P., Sharma, P., & Thakur, N. (2024). Sustainable farming practices and soil health: A pathway to achieving SDGs and future prospects. *Discover Sustainability*, *5*(1), 250.

Song, Y., Li, Z., Ha, J., He, X., Fu, J., Zheng, F., & Li, Z. (2026). Integrated implementation of conservation agriculture enhances crop yield and sustains multiple ecosystem services in the black soil regions. *Agriculture, Ecosystems & Environment*, *411*, 110614.

Sun, S., Zhang, M., Gu, X., He, S., & Tang, L. (2022). Microbial response mechanism of plants and zero valent iron in ecological floating bed: Synchronous nitrogen, phosphorus removal and greenhouse gas emission reduction. *Journal of environmental management*, *324*, 116326.

Teague, R., & Kreuter, U. (2020). Managing grazing to restore soil health, ecosystem function, and ecosystem services. *Frontiers in Sustainable Food Systems*, *4*, 534187.

Vergopolan, N., Xiong, S., Estes, L., Wanders, N., Chaney, N. W., Wood, E. F., ... & Sheffield, J. (2020). Field-scale soil moisture bridges the spatial-scale gap between drought monitoring and agricultural yields. *Hydrology and Earth System Sciences Discussions*, *2020*, 1-31.

Vittori Antisari, L., Ferronato, C., De Feudis, M., Natali, C., Bianchini, G., & Falsone, G. (2021). Soil biochemical indicators and biological fertility in agricultural soils: A case study from northern Italy. *Minerals*, *11*(2), 219.

Von Cossel, M., Scordia, D., Altieri, M., & Gresta, F. (2025). Spotlight on agroecological cropping practices to improve the resilience of farming systems: a qualitative review of meta-analytic studies. *Frontiers in Agronomy*, *7*, 1495846.

Wang, B., Li, W., Xue, N., Xi, R., Wang, Y., Fang, L., ... & Wu, X. (2025). Co-application of biochar and compost enhanced soil carbon sequestration in urban green space. *Frontiers in Microbiology*, *16*, 1707894.

Willoughby, C. M., Topp, C. F., Hallett, P. D., Stockdale, E. A., Walker, R. L., Hilton, A. J., & Watson, C. A. (2023). Soil health metrics reflect yields in long-term cropping system experiments: CM Willoughby et al. *Agronomy for Sustainable Development*, *43*(5), 65.

Xing, Y., Wang, X., & Mustafa, A. (2025). Exploring the link between soil health and crop productivity. *Ecotoxicology and environmental safety*, *289*, 117703.

Yaqoob, N., Farooque, A. A., Shah, S. H. H., Abbas, F., & Hassan, M. (2026). Variable rate irrigation through digital agriculture for sustainable water management: a meta review on current challenges and future directions. *Water Resources Management*, *40*(6), 237.

Yuan, Y., Luo, Y., Wang, H., Ren, Y., Mei, N., Zheng, H., ... & Zheng, J. (2026). A comparative analysis of soil health index calculation methods and minimum dataset construction approaches: a case study from a 40-year tillage experiment. *Ecological Indicators*, *182*, 114609.

Zaghloul, A., Saber, M., Gadow, S., & Awad, F. (2020). Biological indicators for pollution detection in terrestrial and aquatic ecosystems. *Bulletin of the National Research Centre*, *44*(1), 1-11.

Zhang, F., Bei, J., Shi, Q., Wang, Y., & Wu, L. (2024). Research on agricultural machinery services for the purpose of promoting conservation agriculture: an evolutionary game analysis involving farmers, agricultural machinery service organizations and governments. *Agriculture*, *14*(8), 1383.

IEER
Excellence in Education & Research