Policy Research Journal
ISSN (E): 3006-7030 ISSN (P) : 3006-7022 Volume 4, Issue 3, 2026

OPTIMIZING FISH GROWTH AND FEED EFFICIENCY THROUGH SUSTAINABLE NUTRITIONAL STRATEGIES IN AQUACULTURE

Salman[*1], Mahrukh Chaoudary[2], Shanzeel Ahmed[3],

[*1]*Department of Zoology, Division of Science and Technology, University of Education, Township Campus, Lahore 54770, Pakistan*
[2]*Government college women university Faisalabad Bachelor in Zoology*
[3]*Zoology University of education jauharabad campus*
https://orcid.org/0009-0001-8773-3893

[*1]salmanbsf1900322@gmail.com, [2]mahrukhchaoudary62@gmail.com, [3]shanzeelahmed04@gmail.com

DOI: https://doi.org/10.5281/zenodo.23557779

Keywords
sustainable aquaculture; alternative aquafeed ingredients; fishmeal replacement; feed conversion ratio; precision fish farming; circular bioeconomy; immunomodulation

Article History
Received: 07 January 2026
Accepted: 17 March 2026
Published: 31 March 2026

Copyright @Author
Corresponding Author: *
Salman

Abstract

Global aquaculture has surpassed wild-capture fisheries as the primary supplier of aquatic protein, yet its continued expansion is constrained by an overreliance on finite marine-derived ingredients, specifically fishmeal (FM) and fish oil (FO). With feed accounting for 60–70% of operational expenditure in intensive systems, identifying sustainable, economically viable alternatives has become a central research priority. This review synthesizes recent advances in sustainable nutritional strategies aimed at optimizing fish growth performance and feed efficiency in aquaculture. It examines four major alternative ingredient classes terrestrial plant protein sources, rendered animal by-products, insect-derived meals, and single-cell proteins and microalgae evaluating their nutritional profiles, essential amino acid (EAA) compositions, anti-nutritional factors (ANFs), and species-specific inclusion thresholds. Particular attention is given to the physiological, histological, and immunological responses of aquatic species to these ingredients, including intestinal mucosal integrity, gut microbiome modulation, and immune marker expression. The review demonstrates that synergistic multi-ingredient blending, supported by targeted crystalline EAA supplementation, effectively mitigates the limitations of single-source substitution by diluting ANF toxicity and constructing balanced nutrient matrices. Furthermore, the integration of Precision Fish Farming (PFF) technologies computer vision, passive acoustics, and AI-driven multimodal sensor fusion enables dynamic, satiety-based feed delivery that reduces feed conversion ratios (FCR) by 5–30% while minimizing environmental nutrient loading. Finally, sustainability metrics, circular bioeconomy pathways, and the economic bottlenecks impeding commercial scaling are critically assessed. Achieving cost parity and long-term industry adoption will require coordinated advances in processing infrastructure, regulatory standardization, policy incentives, and continued interdisciplinary research.

INTRODUCTION:

The Aquafeed Paradigm Shift

Global aquaculture has established itself as the primary supplier of aquatic protein for human consumption, exceeding wild-capture fisheries production and expanding at a pace that surpasses major terrestrial livestock sectors. This growth has been underpinned by advancements in nutritional science, diet formulation, and intensive rearing systems (Hixson, 2014). However, the historical dependence of aquafeed formulations on marine-derived raw materials specifically fishmeal (FM) and fish oil (FO) presents an environmental and economic bottleneck that threatens the long-term expansion of the industry (Watanabe, 2002).

Fishmeal and fish oil have traditionally served as the gold standard ingredients in aquafeeds, particularly for carnivorous and high-value teleost species, owing to their elevated crude protein levels, balanced essential amino acid (EAA) profiles, superior fatty acid composition (specifically long-chain omega-3 polyunsaturated fatty acids like eicosapentaenoic acid [EPA] and docosahexaenoic acid [DHA]), high digestibility, and high palatability, alongside an absence of anti-nutritional factors (ANFs) (Brum et al., 2025). Nevertheless, global annual production of FM and FO has plateaued at approximately 4.5 million tonnes and 0.9 million tonnes, respectively. Concurrently, feed inputs represent between 60% and 70% of the total operational expenditure in intensive aquaculture enterprises. This economic reality, compounded by price volatility and ecological imperatives to protect forage fisheries, has catalyzed an intensive global research effort aimed at identifying, evaluating, and scaling sustainable alternative feedstuffs (Sarker, 2023).

Optimizing fish growth performance typically measured via Specific Growth Rate (SGR) and Daily Growth Coefficient (DGC) and feed utilization efficiency, represented by the Feed Conversion Ratio (FCR) and Protein Efficiency Ratio (PER), requires maintaining precise nutrient balances (Hasan, 2001). Substituting FM with alternative raw materials is not a simple unit-for-unit replacement; high inclusion rates of single alternative protein sources frequently induce metabolic stress, disrupt intestinal mucosal structural integrity, alter the gut microbiome, downregulate digestive enzyme activities, and compromise immune competence (Roques et al., 2020).

To overcome these physiological limitations, contemporary aquaculture nutrition is shifting from simple single-ingredient substitution models toward strategic, multi-ingredient synergistic blending paired with precision technology. By combining complementary plant protein isolates, rendered animal by-products, insect meals, and single-cell proteins (SCP), formulators can create balanced EAA matrixes that mitigate the negative physiological effects of individual raw materials (Kumar et al., 2026). Concurrently, the integration of Precision Fish Farming (PFF) technologies such as real-time computer vision, passive acoustics, and artificial intelligence (AI) allows feed delivery to be dynamically aligned with instantaneous physiological satiety, directly targeting the physical waste of feeds and minimizing environmental nutrient loads (Goda et al., 2024).



Figure 1: Nanoparticle Incorporation Strategies and Integrated Bio-Functional Mechanisms in Advanced Aquafeed Formulations.

2. Sustainable Alternative Raw Materials and Nutritional Profiles

2.1 Terrestrial Plant Protein Sources and Anti-Nutritional Mitigation

Terrestrial plant protein ingredients (PPIs), including soybean meal (SBM), soy protein concentrate (SPC), pea protein concentrate (PPC), wheat gluten, rapeseed meal, copra meal, and agricultural side-streams such as pumpkin seed cake and cocoa pod husk, represent the most widely used alternatives to FM due to their high production volume, established supply chains, and relative cost efficiency (Quintieri et al., 2023).

Despite these advantages, raw and coarsely processed plant ingredients present distinct nutritional challenges. Most notable is their content of biologically active anti-nutritional factors (ANFs), such as phytic acid, trypsin inhibitors, lectins, saponins, tannins, and non-starch polysaccharides (NSPs). In high-inclusion diets, ANFs reduce protein and lipid digestibility, chelate divalent cations (such as zinc, iron, and magnesium), inhibit endogenous pancreatic enzymes, and induce morphological enteritis (Quevedo-Olaya et al., 2025).

When aquatic species are exposed to raw plant proteins containing high levels of ANFs, the pathophysiological pathway begins with intestinal inflammation characterized by the upregulation of pro-inflammatory cytokines such as NF-κB, IL-1β, and TNF-α. This inflammatory cascade leads to mucosal degradation, including widening of the lamina propria and submucosa, which severely impairs nutrient absorption and induces growth depression (Kaur et al., 2025). High dietary incorporation of unrefined soy products triggers subacute inflammation in the distal intestine of salmonids and marine teleosts, characterized by the widening of the lamina propria and submucosa, loss of enterocyte vacuolation, and altered mucosal architecture. Furthermore, specific plant by-products contain alkaloid compounds such as theobromine in cocoa shell meal (1.3% content) which mechanically limit inclusion rates due to metabolic toxicity (Petrusán et al., 2016).

Plant proteins also exhibit inherent EAA imbalances, typically displaying marked deficiencies in methionine, lysine, threonine, and tryptophan compared to the physiological requirements of teleost fish. To circumvent these limitations, processing techniques such as thermomechanical extrusion, aqueous ethanol extraction (yielding concentrates with >65% crude protein), microbial fermentation, and enzymatic hydrolysis are deployed to reduce ANF concentrations (Ahmed et al., 2026). When supported by targeted supplementation with crystalline EAAs (e.g., L-methionine and L-lysine), plant protein concentrates can replace up to 40–100% of FM in select omnivorous species like Nile tilapia (*Oreochromis niloticus*) and milkfish (*Chanos chanos*), and up to 25–50% in carnivorous species without severely depressing growth (Wang et al., 2026).

2.2 Rendered Terrestrial Animal By-Products

Rendered terrestrial animal by-products including poultry by-product meal (PBM), feather meal (FEM), blood meal (BM), and meat and bone meal (MBM) represent high-protein raw materials that support circular economy initiatives within animal agriculture. These ingredients are naturally free of plant-derived ANFs, possess favorable crude lipid levels, display superior palatability, and provide high crude protein fractions (ranging from 55% in MBM up to 80–90% in BM and hydrolyzed FEM) (Woodgate et al., 2023).

Studies evaluating animal by-products have demonstrated that complete (100%) replacement of FM can be achieved using BM alone or in synergistic combination with PBM and FEM in species like milkfish and tilapia without impairing overall weight gain. However, unbalanced inclusion of rendered meals can introduce nutritional bottlenecks (Bureau & Meeker, 2010). Blood meal is rich in lysine but severely deficient in isoleucine and methionine, whereas feather meal exhibits low protein digestibility unless extensively hydrolyzed to disrupt keratin disulphide bonds. Additionally, excessive incorporation of rendered animal meals (>80% MBM inclusion) has been shown to downregulate intestinal digestive enzymes (trypsin and lipase activity) and depress endogenous antioxidant capacities, marked by significant decreases in superoxide dismutase (SOD) activity and reduced glutathione (GSH) content (Rodehutscord et al., 2002).

2.3 Insect-Derived Protein Meals

Insect meals have emerged as functional, high-value alternative protein sources, with primary commercial focus centered on Black Soldier Fly Larvae Meal (BSFLM / BSFM; *Hermetia illucens*), Yellow Mealworm (*Tenebrio molitor*), and Housefly Larvae (*Musca domestica*). Insects possess crude protein levels ranging from 40% to 70% on a dry matter (DM) basis and feature favorable EAA profiles that closely align with marine proteins, particularly regarding lysine, leucine, and valine content (Lamsal et al., 2019).

The physiological impact of insect meal inclusion follows a dual dose-dependent response. At moderate inclusion levels, dietary chitin and lauric acid modulate the gut microbiome by encouraging the proliferation of beneficial bacteria such as *Lactobacillus* species, while natural bioactive peptides enhance non-specific immune responses, including lysozyme, immunoglobulin M (IgM), and complement component 4 (C4) activity (Vale-Hagan et al., 2023). However, when dietary inclusion exceeds species-specific thresholds (typically 30–50% replacement), the high chitin content increases structural indigestibility, ultimately depressing feed conversion ratios and growth performance (Lisboa et al., 2024).

In addition to macronutrients, insect meals contain naturally occurring functional compounds, including lauric acid (a medium-chain fatty acid with innate antimicrobial properties), bioactive antimicrobial peptides (AMPs), and chitin. At moderate dietary inclusion rates (typically 10% to 20% total diet, replacing 20% to 33% of FM), BSFLM upregulates the expression of immune-related genes, increases serum lysozyme, IgM, and C4 levels, and enriches beneficial intestinal microbiota such as *Lactobacillus* species (Gasco et al., 2019). In crustacean aquaculture, such as Pacific white

shrimp (*Litopenaeus vannamei*), replacing up to 20% of FM with defatted insect meal enhances resistance against bacterial pathogens like *Vibrio parahaemolyticus* (the causative agent of Acute Hepatopancreatic Necrosis Disease, AHPND) (Hermans et al., 2021).

However, high dietary inclusions of insect meal (>30% to 50% FM replacement, or >10% total inclusion in strict carnivores like juvenile gilthead sea bream *Sparus aurata*) frequently cause growth depression and elevated FCR. This effect is attributed to the non-digestible structural chitin matrix binding dietary nutrients, alongside lipid imbalances characterized by high saturated fatty acid (SFA) profiles and an absence of EPA and DHA (Hobbi et al., 2022).

2.4 Single-Cell Proteins (SCP) and Microalgae

Single-Cell Proteins (SCP) derived from processed, non-viable microbial biomass including methane-utilizing bacteria (*Methylococcus capsulatus*), yeasts (*Saccharomyces cerevisiae*), and microalgae (*Spirulina platensis*, *Schizochytrium* sp., *Chlorella* sp.) represent scalable, land-independent nutritional solutions. Bacteria and yeasts contain crude protein fractions ranging between 50–80% and 45–55% DM, respectively, featuring EAA profiles rich in lysine, threonine, and tryptophan that closely match FM (Ritala et al., 2017).

Microalgal species present distinct physiological and nutritional advantages. *Spirulina platensis* lacks hemicellulose, lignin, and cellulosic cell walls, rendering its protein and polysaccharide matrix highly digestible compared to green microalgae like *Chlorella*, which possess recalcitrant cellulosic cell walls requiring mechanical disruption (e.g., bead milling) for optimal nutrient access. Heterotrophic microalgae, such as *Schizochytrium* sp., function primarily as Single-Cell Oils (SCO), accumulating lipid fractions exceeding 200 g/kg DM that are exceptionally rich in DHA (Bratosin et al., 2021).

Replacing FM and FO with combinations of bacterial SCP and microalgal oil at 15% to 50% substitution levels maintains or enhances Specific Growth Rate (SGR), improves Protein Efficiency Ratio (PER), elevates lysozyme activity, and spares *n*-3 long-chain polyunsaturated fatty acids (*n*-3 LC-PUFA) in fish muscle tissue. However, inclusion rates exceeding 50% bacterial or yeast SCP can lead to reduced feed intake due to palatability shifts and potential nucleic acid accumulation, requiring careful dietary calibration (Barka & Blecker, 2016).

Table 1: Comparative Nutritional Composition, Inclusion Thresholds, and Functional Impacts of Alternative Aquafeed Ingredients

Ingredient Class	Representative Ingredients	Crude Protein Range (% DM)	Key EAA Deficiencies / ANFs	Recommended Inclusion (FM Replacement %)	Main Physiological & Immunological Effects
Plant Protein Concentrates [cite: 3, 4, 5, 8, 14]	Soy Protein Concentrate (SPC), Pea Protein Concentrate (PPC)	60 – 75%	Deficient in Methionine, Lysine. ANFs: Phytic acid, lectins, saponins	25 – 50% (Carnivores)	

40 – 100% (Omnivores) | Risk of distal enteritis, widened lamina propria; requires crystalline EAA supplementation. | | **Rendered Animal By-Products** | Poultry By-Product Meal (PBM), Blood Meal (BM), Feather Meal (FEM) | 55 – 90% | BM: Deficient in Isoleucine. FEM: Variable EAA digestibility | 50 – 100% (Partial or Total Combinations) | High palatability; excessive MBM (>80%) depresses digestive enzymes and antioxidant markers (SOD, GSH) (Hossain et al., 2024). | | **Insect-Derived Meals** | Black Soldier Fly Larva (*H. illucens*), Mealworm (*T. molitor*) | 40

– 70% | Lacks EPA/DHA; High structural chitin content | 10 – 33% (Up to 20% total diet inclusion) | Promotes gut *Lactobacillus*, boosts IgM, lysozyme, and C4; chitin depresses FCR at >30–50% inclusion. | | **Single-Cell Proteins (SCP)** | Bacterial Protein (*M. capsulatus*), Yeast (*S. cerevisiae*) | 45 – 80% | Favorable EAA profile; high nucleic acid content | 15 – 50% | Enhances growth, PER, and gut architecture; high inclusion (>50%) may reduce palatability (Jiang et al., 2026). | | **Microalgae & Macroalgae** | *Spirulina platensis*, *Schizochytrium* sp., Seaweeds | 30 – 70% | Varied protein; *Chlorella* cell walls require disruption | 5 – 25% (Proteins & Lipid SCOs) | *Spirulina* is cell-wall-free and highly digestible; *Schizochytrium* provides EPA/DHA and boosts antioxidant capacity (Nazir et al., 2026). |

3. Synergistic Multi-Ingredient Formulation and Health Responses

3.1 Advanced Multi-Ingredient Blending Concepts

Single-ingredient substitution strategies are inherently constrained by the unique nutritional imbalances of individual raw materials. Plant proteins introduce ANFs and lack sulfur-containing amino acids; animal by-products exhibit EAA variability; insect meals lack long-chain n-3 fatty acids and carry non-digestible chitin fractions; and single-cell proteins can be cost-prohibitive at high inclusion levels (Hao et al., 2025).

The paradigm of modern aquafeed formulation relies on synergistic multi-ingredient blending. The complementary matrix approach integrates plant protein concentrates (which provide high crude protein but are low in methionine), insect larval meals (rich in lysine and chitin), and single-cell proteins or microalgae meals (rich in methionine, EPA/DHA, and antimicrobial peptides). Combining these diverse feedstuffs yields a balanced essential amino acid and fatty acid matrix, dilutes single-source anti-nutritional factor toxicity, and preserves mucosal structural integrity and feed conversion efficiency (Hassan et al., 2024).

By combining complementary protein and lipid matrixes for instance, pairing methionine-deficient soy concentrates with methionine-rich microalgae or bacterial SCP, and supplementing defatted insect meal with DHA-rich *Schizochytrium* oil formulators can construct balanced nutrient profiles (Jacobo-Velázquez, 2024). This complementary approach mitigates the toxicological thresholds of individual ANFs, optimizes total nutrient digestibility, and eliminates the growth depression associated with single-source replacement strategies (Nilsson et al., 2023).

3.2 Intestinal Histology, Digestive Physiology, and Immunomodulation

The inclusion of alternative raw materials directly modulates the mucosal morphology, enzymatic activity, and immunological pathways of the aquatic host. Ingestion of unrefined plant meal downregulates hypoxia-inducible factor 1 (HIF-1) and upregulates key pro-inflammatory cytokines (Andrade et al., 2025). Including nuclear factor kappa B (NF-κB p65), tumor necrosis factor-alpha (TNF-α), interleukin 1-beta (IL-1β), and interleukin 10 (IL-10). This inflammatory cascade induces structural degradation of the intestine, characterized by an increase in lamina propria width (LPW) and submucosa width (SMW) (Valeur et al., 2004).

Conversely, incorporating functional alternative protein sources restores mucosal homeostasis. Inclusion of fermented plant meals, insect larvae meals, or bacterial SCP upregulates anti-inflammatory pathways (transforming growth factor-beta, TGF-β), suppresses pro-inflammatory cascades, and enhances brush-border enzyme activities (trypsin, lipase, and alkaline phosphatase) (Arrieta et al., 2011). From an immunological perspective, moderate inclusions of insect meal (e.g., up to 200 g/kg BSFLM) and single-cell ingredients significantly elevate systemic immune markers, including serum lysozyme, nitric oxide production, and total IgM (Koon & Pothoulakis, 2006).

Furthermore, these ingredients exert positive prebiotic effects on the gut microbiome; for

example, BSFLM and microalgal polysaccharides increase the relative abundance of beneficial lactic acid bacteria (*Lactobacillus* spp.), which competitively exclude opportunistic pathogens (e.g., *Aeromonas hydrophila*, *Aeromonas veronii*, and *Vibrio* species) and strengthen intestinal barrier integrity (Siddiqui & Cresci, 2021).

Table 2: Key Physiological, Histological, and Immunomodulatory Effects of Sustainable Aquafeed Formulations Across Species

Species	Formulated Diet / Ingredient Substitution	Inclusion / Substitution Threshold	Physiological & Histological Response	Immunomodulatory & Health Outcome
Koi Carp (*Cyprinus carpio* var. *koi*)	Black Soldier Fly Larva Meal (BSFLM) replacing FM	Up to 200 g/kg (20% inclusion)	Higher Weight Gain and SGR; maintained optimal FCR and 100% survival	Upregulated immune-related gene expression; enhanced innate disease resistance.
Nile Tilapia (*Oreochromis niloticus*)	Pumpkin Seed Cake / Copra Meal replacing Soybean Meal	35% replacement of SBM (10.5% diet inclusion)	Significant increase in final weight, SGR (up to 4.89%/day), and PER (2.69)	Improved nutrient utilization and feed efficiency.
European Sea Bass (*Dicentrarchus labrax*)	Single-Cell Ingredients (SCP + Microalgae) replacing FM/FO	15% combined inclusion (9.4% SCP, 4.7% yeast, 1% algae)	Improved anterior/posterior gut mucosal structure; higher nutrient digestibility	Significant increase in serum lysozyme activity; maintained growth performance.
Gilthead Sea Bream (*Sparus aurata*)	*Methylococcus capsulatus* SCP vs. BSF Insect Meal	66% FM replacement by SCP; 33% FM replacement by BSF	SCP supported full growth performance; BSF at 10% inclusion depressed growth	SCP increased fillet concentrations of 20:5*n*-3 (EPA) and 22:6*n*-3 (DHA).
Pacific White Shrimp (*L. vannamei*)	Defatted Insect Larvae Meal replacing FM	Up to 20% FM replacement	Maintained normal growth rate and feed conversion; higher inclusion (>30%) caused degenerative effects	Modulated intestinal microbiome; enhanced disease resistance against AHPND.

4. Precision Aquaculture and Smart Feeding Technologies

4.1 Integration of Precision Fish Farming Paradigms

While optimizing diet chemistry is essential, feed efficiency is equally dependent on delivery dynamics. Feed constitutes up to 70% of farm operational costs; overfeeding leads to unconsumed feed pellets sinking to the benthic zone, degrading water quality, driving up FCR, and inducing environmental eutrophication (Lin et al., 2026). Precision Fish Farming (PFF) also designated as Smart Aquaculture or Fish Farming 5.0 applies Internet of Things (IoT) frameworks, sensor networks, edge computing, and artificial intelligence (AI) to transform feeding management into a data-driven, closed-loop process (Føre et al., 2023).

In a precision feeding architecture, input streams from visual tracking cameras, infrared/LiDAR

sensors, passive acoustic hydrophones, and environmental water quality sensors feed directly into edge and cloud AI analytics platforms. Algorithms such as DMRI-Net and ResNet-CBAM execute multimodal sensor fusion via Bayesian inference to determine dynamic satiety endpoints, continuously regulating smart automated feeders to reduce feed conversion ratios by 5% to 30% (D'Agaro, 2025).

Continuous environmental monitoring tracking dissolved oxygen (DO), temperature, pH, and ammonia is paired with real-time bio-behavioral sensing. Data collected from fish tanks, raceways, or sea cages is transmitted via wireless sensor networks (WSNs) using protocols such as LoRaWAN to cloud analytics engines (Yavuzcan & Mansour, 2026). Automated Decision Support Systems (DSS) process these data streams to dynamically regulate feed distribution rates based on calculated biological demand rather than static feeding tables (Zhang et al., 2025).



Figure 2: Schematic Representation of Nitrogen Bioconversion and Recirculating Nutrient Dynamics in an Integrated Aquaponics Framework.

4.2 Sensing Modalities and Algorithmic Satiety Detection

Intelligent feeding systems utilize three primary non-invasive sensing modalities: computer vision, hydro-acoustics, and biological logging (Irshad et al., 2022).

Computer Vision and Optical Sensing

Optical camera systems (monocular, stereo vision, near-infrared [NIR], and Light Detection and Ranging [LiDAR]) acquire high-resolution imagery to analyze fish spatial distribution, swimming speed, biomass, and surface feeding activity (Tarabanis et al., 1995). Mono and stereo systems calculate 3D spatial coordinates to track individual fish morphology and biomass with margins of error under 3–5%. In outdoor or turbid environments where sub-surface optical clarity is compromised, surface vision systems rely on deep learning convolutional neural networks (CNNs) to analyze feeding turbulence (Yuan et al., 2023).

For example, specialized deep learning architectures (such as DMRI-Net, R(2+1)D deep models, and ResNet variants augmented with Convolutional Block Attention Modules [CBAM]) evaluate the intensity of water surface

waves generated during feeding, achieving classification accuracies between 93.2% and 99.72% for satiety estimation (Peters, 2017).

Hydro-Acoustic Technologies

In deep sea-cages or high-turbidity land-based Recirculating Aquaculture Systems (RAS) where optical systems suffer from light attenuation, passive hydro-acoustic sensors provide a continuous monitoring alternative (Li et al., 2024). Underwater hydrophones capture the acoustic signatures generated during feed ingestion, specifically the distinct high-frequency sound pulses created by pellet mastication and crunching. Algorithms quantify the frequency and amplitude of these mastication signals, establishing a direct correlation with herd satiety. When sound pulse frequency drops below designated thresholds, automated feed dispensers immediately halt delivery (Shabangu et al., 2014).

Multimodal Sensor Fusion

The most advanced precision feeding platforms utilize multimodal data fusion. By integrating optical surface wave analytics, underwater acoustic mastication signals, and implantable bio-logger metrics (measuring real-time electromyograms [EMG] or acceleration rates), AI frameworks apply Bayesian change-point detection algorithms to determine exact satiety endpoints (Lahat et al., 2015).

Field implementations of these integrated technologies demonstrate operational improvements: feed conversion ratios (FCR) decrease by approximately 5% to 30%, mortality drops by up to 40%, water use efficiency increases by up to 50%, and total biomass yield is optimized (Chung et al., 2019).

Table 3: Technical Comparison of Sensing Technologies in Precision Aquaculture Feeding Systems

Sensing Modality	Operational Mechanism	Primary Aquaculture Application	Classification / Measurement Accuracy	Major Operational Strengths & Constraints
Monocular & Stereo Vision [cite: 12, 22]	Dual optical cameras capturing overlapping spatial coordinates for 3D measurement	Non-invasive biomass estimation, morphological profiling, and spatial tracking	High spatial precision; biomass error margin within 3–5%	Strengths: Detailed individual tracking.

5. Sustainability Metrics, Circular Bioeconomy, and Economic Feasibility

5.1 Re-evaluating Traditional Sustainability Metrics

Assessing the ecological sustainability of aquafeed formulations has historically relied on two standard metrics: the Fish-In:Fish-Out (FIFO) ratio and the Forage Fish Dependency Ratio (FFDR) (Yang et al., 2026). The FIFO metric quantifies the mass of wild forage fish required to produce a unit mass of farmed fish, calculated as:

$$\text{FIFO} = \frac{\text{Dietary FM Inclusion} + \text{Dietary FO Inclusion}}{\text{Yield of FM from Wild Fish} + \text{Yield of FO from Wild Fish}} \times \text{FCR}$$

While FIFO and FFDR provided early benchmarks for reducing wild fish inputs, contemporary life-cycle resource economists highlight their analytical limitations (Latinović et al., 2026). These metrics fail to differentiate between poorly managed and certified sustainable forage fisheries, nor do they account for the utilization of fishery processing trimmings (by-products). Furthermore, FIFO ignores ecological value conversion such as converting low-cost small pelagics like anchoveta into high-value protein and overlooks broader environmental impacts like terrestrial land utilization, freshwater consumption, global warming potential (GWP), and eutrophication potential (Clark et al., 2022).

Consequently, sustainability assessments are adopting comprehensive Life Cycle Assessment (LCA) frameworks. LCAs evaluate cradle-to-gate impact parameters across the complete ingredient supply chain, including land-use transformation for plant crops, energy inputs for microbial single-cell drying, and carbon footprints associated with insect rearing substrates (Binder, 2024).

5.2 Circular Bioeconomy and Industrial Waste-to-Feed Pathways

The long-term viability of alternative aquafeed ingredients depends on alignment with circular bioeconomy principles. The circular bioeconomy pipeline converts organic side-streams and industrial carbon dioxide emissions into high-protein insect biomass and single-cell microbial meals through gas fermentation and bioconversion (Dhiman et al., 2026). These bio-meals are blended into sustainable aquafeed formulations to replace finite marine fishmeal and fish oil, delivering optimized fish growth rates while minimizing land, water, and carbon footprints (Nath et al., 2023).

Novel ingredients valorize industrial waste streams, transforming low-value side-streams into high-value nutrients. Insect farming utilizes municipal food waste, agricultural residues, and distillery grains to produce high-protein larvae biomass. Similarly, gas-fermentation bioprocesses capture industrial carbon emissions (such as CO_2 or methane) to cultivate single-cell bacterial proteins (*M. capsulatus*), bypassing reliance on arable land and fresh water (Bowler et al., 2026).

5.3 Economic Realities and Scaling Bottlenecks

The adoption of alternative feedstuffs by commercial feed mills is dictated by ingredient price parity, supply chain stability, and regulatory compliance. While standard plant meals offer lower costs, premium concentrates, insect meals, and single-cell oils present varied cost structures (Chang & Andreoni, 2020).

Table 4: Economic Profile and Commercialization Trajectory of Primary Aquafeed Raw Materials

Ingredient Category	Estimated Market Price (USD / Metric Tonne)	Supply Chain Scalability Status	Primary Economic & Technological Bottlenecks
Conventional Fishmeal (FM)	~$1,800 / MT (High market volatility)	Globally established; volume capped at ~4.5M tonnes/year	Resource constraints, forage fishery limits, and high price sensitivity.
Soy Protein Concentrate (SPC)	~$700 / MT	Highly scaled; globally distributed commercial supply	Competition with terrestrial livestock feed and human food sectors.
Defatted Insect Meal (BSFLM)	~$2,500 / MT	Emerging; localized production facilities expanding	High production costs, fragmented production volumes, substrate regulatory limits.
Heterotrophic Microalgae Meal / Oil	~$5,000 / MT	Industrial scale emerging; expanding capacity	High energy expenditure for cell drying and mechanical cell-wall disruption.

Alternative protein meals like defatted insect larvae and microalgal oils currently carry higher unit prices than standard FM or plant isolates. Achieving cost parity requires commercial scaling, public policy support, capital investment incentives for processing infrastructure, and regulatory standardization regarding waste substrates used for insect and microbial cultivation (Hasan & Shipton, 2021).

REFERENCES

Ahmed, N., Sofi, S. A., & Sheikh, M. A. (Eds.). (2026). *Pulse Proteins: Food Product Formulation*. CRC Press.

Andrade, M. E. R., Araújo, R. S., de Barros, P. A. V., Soares, A. D. N., Abrantes, F. A., de Vasconcelos Generoso, S., ... & Cardoso, V. N. (2015). The role of immunomodulators on intestinal barrier homeostasis in experimental models. *Clinical nutrition*, *34*(6), 1080-1087.

Arrieta, M. C., Meddings, J., & Field, C. J. (2011). The immunomodulatory effects of dietary fiber and prebiotics in the gastrointestinal tract. *Non digestible Carbohydrates and Digestive Health*, 37-77.

Barka, A., & Blecker, C. (2016). Microalgae as a potential source of single-cell proteins. A review. *Biotechnologie, Agronomie, Société et Environnement*, *20*(3).

Binder, P. M. (2024). Mutiple valorization pathways to recover high value resources from dairy wastewater and beyond.

Bowler, A. L., Pennells, J., Knoerzer, K., Eastham, L., Jarvis, A., & Watson, N. J. (2026). Fermentelligence–Can AI-optimised fermentation of agri-food side streams deliver the sustainable protein promise?. *Current Opinion in Green and Sustainable Chemistry*, 101091.

Bratosin, B. C., Darjan, S., & Vodnar, D. C. (2021). Single cell protein: A potential substitute in human and animal nutrition. *Sustainability*, *13*(16), 9284.

Brum, A., Magnotti, C., Tsuzuki, M. Y., Sousa, E. M. D. O., Mouriño, J. L. P., Martins, M. L., ... & Owatari, M. S. (2025). Pivotal roles of fish nutrition and feeding: recent advances and future outlook for Brazilian fish farming. *Fishes*, *10*(2), 47.

Bureau, D. P., & Meeker, D. L. (2010). Terrestrial animal fats. *Fish oil replacement and alternative lipid sources in aquaculture feeds. CRC Press, Boca Raton, FL*, 245-266.

Chang, H. J., & Andreoni, A. (2020). Industrial policy in the 21st century. *Development and change*, *51*(2), 324-351.

Chung, S., Lim, J., Noh, K. J., Kim, G., & Jeong, H. (2019). Sensor data acquisition and multimodal sensor fusion for human activity recognition using deep learning. *Sensors*, *19*(7), 1716.

Clark, J. H., Jardine, A., Matharu, A. S., & Stevens, C. V. (2022). *Greener Organic Transformations*. Royal Society of Chemistry.

D'Agaro, E. (2025). Fish farming 5.0: Advanced tools for a smart aquaculture management. *Applied Sciences*, *15*, 1-24.

Dhiman, S., Mukherjee, G., Kumar, A., & Krishan, B. (2026). Alternative proteins from waste streams: a circular economy roadmap for sustainable food systems. *Sustainable Food Technology*, *4*(3), 2266-2294.

Føre, M., Alver, M. O., Frank, K., & Alfredsen, J. A. (2023). Advanced technology in aquaculture–smart feeding in marine fish farms. In *Smart livestock nutrition* (pp. 227-268). Cham: Springer International Publishing.

Gasco, L., Biasato, I., Dabbou, S., Schiavone, A., & Gai, F. (2019). Animals fed insect-based diets: State-of-the-art on digestibility, performance and product quality. *Animals*, *9*(4), 170.

Goda, A. M. S., Aboseif, A. M., Taha, M. K., Mohammady, E. Y., Aboushabana, N. M., Nazmi, H. M., ... & Ashour, M. (2024). Optimizing nutrient utilization, hydraulic loading rate, and feed conversion ratios through freshwater IMTA-aquaponic and hydroponic systems as an environmentally sustainable aquaculture concept. *Scientific Reports*, *14*(1), 14878.

Hao, M., Tan, X., Liu, K., & Xin, N. (2025). Nanoencapsulation of nutraceuticals: Enhancing stability and bioavailability in functional foods. *Frontiers in Nutrition*, *12*, 1746176.

Hasan, M. R. (2001, February). Nutrition and feeding for sustainable aquaculture development in the third millennium. In *Aquaculture in the third millennium. Technical proceedings of the conference on aquaculture in the third millennium* (pp. 193-219). Thailand: Bangkok.

Hasan, M. R., & Shipton, T. A. (2021). Aquafeed value chain analysis of striped catfish in Vietnam. *Aquaculture*, *541*, 736798.

Hassan, I., Gani, A., & Mudabir, S. (2024). Development and characterization of 3D printed fish gelatin and vegetable blends as nutritious meal for dysphagic patients. *Food Hydrocolloids*, *157*, 110478.

Hermans, W. J., Senden, J. M., Churchward-Venne, T. A., Paulussen, K. J., Fuchs, C. J., Smeets, J. S., ... & van Loon, L. J. (2021). Insects are a viable protein source for human consumption: from insect protein digestion to postprandial muscle protein synthesis in vivo in humans: a double-blind randomized trial. *The American journal of clinical nutrition*, *114*(3), 934-944.

Hixson, S. M. (2014). Fish nutrition and current issues in aquaculture: the balance in providing safe and nutritious seafood, in an environmentally sustainable manner.

Hobbi, P., Bekhit, A. E. D. A., Debaste, F., Lei, N., & Shavandi, A. (2022). Insect-derived protein as food and feed. In *Alternative proteins* (pp. 85-132). CRC Press.

Hossain, M. S., Small, B. C., Kumar, V., & Hardy, R. (2024). Utilization of functional feed additives to produce cost-effective, ecofriendly aquafeeds high in plant-based ingredients. *Reviews in Aquaculture*, *16*(1), 121-153.

Irshad, M. T., Nisar, M. A., Huang, X., Hartz, J., Flak, O., Li, F., ... & Grzegorzek, M. (2022). SenseHunger: Machine learning approach to hunger detection using wearable sensors. *Sensors*, *22*(20), 7711.

Jacobo-Velázquez, D. A. (2025). Functional foods for cholesterol management: A review of the mechanisms, efficacy, and a novel cholesterol-lowering capacity index. *Nutrients*, *17*(16), 2648.

Jiang, W., Wu, P., Zhang, H., Ma, Y., Liu, Y., Zhou, X., & Feng, L. (2026). Functional additives in fishmeal-replaced aquafeeds: precision strategies for sustainable aquaculture. *Journal of Animal Science and Biotechnology*, *17*(1), 146.

Kaur, S. P., Sagar, N. A., & Rani, N. (2025). Alternative proteins: Innovations in sources, processing, and consumption. *Frontiers in Sustainable Food Systems*, *9*, 1641712.

Koon, H. W., & Pothoulakis, C. (2006). Immunomodulatory properties of substance P: the gastrointestinal system as a model. *Annals of the New York Academy of Sciences*, *1088*(1), 23-40.

Kumar, S., Banik, A., Dubey, M. K., Srivastava, P. P., & Sandor, Z. J. (2026). Early nutritional programing: unlocking the potential of fish for sustainable aquaculture. *Aquaculture Nutrition*, *2026*(1), 3380126.

Kumar, S., Banik, A., Dubey, M. K., Srivastava, P. P., & Sandor, Z. J. (2026). Early nutritional programing: unlocking the potential of fish for sustainable aquaculture. *Aquaculture Nutrition*, *2026*(1), 3380126.

Lahat, D., Adali, T., & Jutten, C. (2015). Multimodal data fusion: an overview of methods, challenges, and prospects. *Proceedings of the IEEE*, *103*(9), 1449-1477.

Lamsal, B., Wang, H., Pinsirodom, P., & Dossey, A. T. (2019). Applications of insect-derived protein ingredients in food and feed industry. *Journal of the American Oil Chemists' Society*, *96*(2), 105-123.

Latinović, L., Marković, S., & Tomašević, V. (2026). Beyond waste valorization: Glycerol-based metalworking fluids as Hephaestus for the circular economy and sustainability transitions. *Challenges in Sustainability*, *14*(1), 169-192.

Li, D., Du, Z., Wang, Q., Wang, J., & Du, L. (2024). Recent advances in acoustic technology for aquaculture: A review. *Reviews in Aquaculture*, *16*(1), 357-381.

Lin, K., Zhang, S., Wang, W., Dong, T., Ding, W., Geng, C., ... & Hu, H. (2026). Advancing Non-Contact Fish Phenotyping via Optical Sensing: Toward Precision Aquaculture. *Reviews in Aquaculture*, *18*(1), e70123.

Lisboa, H. M., Nascimento, A., Arruda, A., Sarinho, A., Lima, J., Batista, L., ... & Andrade, R. (2024). Unlocking the potential of insect-based proteins: Sustainable solutions for global food security and nutrition. *Foods*, *13*(12), 1846.

Nath, P. C., Ojha, A., Debnath, S., Sharma, M., Nayak, P. K., Sridhar, K., & Inbaraj, B. S. (2023). Valorization of food waste as animal feed: a step towards sustainable food waste management and circular bioeconomy. *Animals*, *13*(8), 1366.

Nazir, M. I., Gargotra, P., Asimi, O., Bhat, I. A., Iqbal, G., & Aakash, M. (2026). Omics-guided aquafeed innovation: raw ingredients to functional nutrition. *Frontiers in Fish Science*, *4*, 1856912.

Nilsson, M. I., May, L., Roik, L. J., Fuda, M. R., Luo, A., Hettinga, B. P., ... & Tarnopolsky, M. A. (2023). A multi-ingredient supplement protects against obesity and infertility in western diet-fed mice. *Nutrients*, *15*(3), 611.

Peters, J. F. (2017). *Foundations of computer vision* (pp. 3-17). New York: Springer International Publishing.

Petrusán, J. I., Rawel, H., & Huschek, G. (2016). Protein-rich vegetal sources and trends in human nutrition: A review. *Curr. Top. Pept. Protein Res*, *17*, 1-19.

Quevedo-Olaya, J. L., Schmiele, M., & Correa, M. J. (2025). Potential of Andean grains as substitutes for animal proteins in vegetarian and vegan diets: A nutritional and functional analysis. *Foods*, *14*(17), 2987.

Quintieri, L., Nitride, C., De Angelis, E., Lamonaca, A., Pilolli, R., Russo, F., & Monaci, L. (2023). Alternative protein sources and novel foods: benefits, food applications and safety issues. *Nutrients*, *15*(6), 1509.

Ritala, A., Häkkinen, S. T., Toivari, M., & Wiebe, M. G. (2017). Single cell protein state-of-the-art, industrial landscape and patents 2001–2016. *Frontiers in microbiology*, *8*, 2009.

Rodehutscord, M., Abel, H. J., Friedt, W., Wenk, C., Flachowsky, G., Ahlgrimm, H. J., ... & Breves, G. (2002). Consequences of the ban of by-products from terrestrial animals in livestock feeding in Germany and the European Union: alternatives, nutrient and energy cycles, plant production, and economic aspects. *Archives of Animal Nutrition*, *56*(2), 67-91.

Roques, S., Deborde, C., Richard, N., Skiba-Cassy, S., Moing, A., & Fauconneau, B. (2020). Metabolomics and fish nutrition: a review in the context of sustainable feed development. *Reviews in Aquaculture*, *12*(1), 261-282.

Sarker, P. K. (2023). Microorganisms in fish feeds, technological innovations, and key strategies for sustainable aquaculture. *Microorganisms*, *11*(2), 439.

Shabangu, F. W., Coetzee, J. C., Hampton, I., Kerwath, S. E., de Wet, W. M., & Lezama-Ochoa, A. (2014). Hydro-acoustic technology and its application to marine science in South Africa. *Marine and Maritime Sectors*, 122.

Siddiqui, M. T., & Cresci, G. A. (2021). The immunomodulatory functions of butyrate. *Journal of inflammation research*, 6025-6041.

Tarabanis, K. A., Allen, P. K., & Tsai, R. Y. (1995). A survey of sensor planning in computer vision. *IEEE transactions on Robotics and Automation*, *11*(1), 86-104.

Vale-Hagan, W., Singhal, S., Grigoletto, I., Totaro-Fila, C., Theodoridou, K., & Koidis, A. (2023). Edible insects in mixed-sourced protein meals for animal feed and food: An EU focus. *Food and Humanity*, *1*, 1180-1187.

Valeur, N., Engel, P., Carbajal, N., Connolly, E., & Ladefoged, K. (2004). Colonization and immunomodulation by Lactobacillus reuteri ATCC 55730 in the human gastrointestinal tract. *Applied and Environmental Microbiology*, *70*(2), 1176-1181.

Wang, Z., Liu, H., Rizvi, S. S., Zhou, L., & Ren, F. (Eds.). (2026). *Plant Proteins, Bioactive Peptides, and Healthy Foods: Nutritional Properties, Functional Applications, and Processing Innovations*. Academic Press.

Watanabe, T. A. K. E. S. H. I. (2002). Strategies for further development of aquatic feeds. *Fisheries science*, *68*(2), 242-252.

Woodgate, S. L. (2023). Meat industry by-products: a bio-refinery approach to the production of safe, value added products for sustainable agriculture applications. *Frontiers in Animal Science*, *4*, 1259200.

Yang, H., Zhang, Q., Diao, H., & Xia, S. (2026). A Bibliometric Analysis and Comprehensive Review on the Valorization of Areca catechu Wastes: From Environmental Burden to Sustainable Resources. *ACS Sustainable Resource Management*, *3*(5), 1695-1714.

Yavuzcan, H., & Mansour, A. T. (2026). Technological innovations are driving precision aquaculture towards a future of transparent, automated, and sustainably efficient production. *Animal Reports*, *2*(1), 16-34.

Yuan, S., Ma, C., Fetaya, E., Mueller, T., Naveh, D., Zhang, F., & Xia, F. (2023). Geometric deep optical sensing. *Science*, *379*(6637), eade1220.

Zhang, S., Li, D., Zhao, J., Yao, M., Chen, Y., & Wang, H. (2025). A comprehensive review of fish feeding behavior analysis in aquaculture: tasks, techniques, and applications. *arXiv preprint arXiv:2502.15311*.

IEER
Institute for Excellence in Education & Research