Policy Research Journal
ISSN (E): 3006-7030 ISSN (P) : 3006-7022 Volume 4, Issue 3, 2026

CLIMATE-RESILIENT NURSERY OPERATIONS, COMBINING SMART PROPAGATION WITH MACHINE LEARNING FOR SEEDLING GROWTH PREDICTION

Ambreen Zain[*1], Tauqeer Nasir[2], Talha[3]

[*1]*Institute of horticulture sciences, university of agriculture Faisalabad, pakistan.*
[2]*Department of Agriculture Extension Allama labal Open University, Islamabad*
[3]*Department of Botany, University of Makran, Panjgur*

[*1]ambreenzain2023@gmail.com, [2]tauqeernasir888@gmail.com, [3]talha@uomp.edu.pk

DOI: https://doi.org/10.5281/zenodo.23611787

Keywords
Climate resilience, smart nursery, IoT sensor networks, precision irrigation, micropropagation bioreactors, LED photomorphogenesis, predictive growth modeling, digital twins

Article History
Received: 07 January 2026
Accepted: 17 March 2026
Published: 31 March 2026

Copyright @Author
Corresponding Author: *
Ambreen zain

Abstract

Global agriculture and forestry face escalating vulnerabilities from climate volatility, with early-stage plant propagation representing the most environmentally sensitive phase of the crop lifecycle. Traditional nursery operations relying on empirical observation and calendar-based practices suffer from substantial water wastage, non-uniform growth, and high sapling mortality. This review synthesizes modern smart propagation technologies that are transforming climate-resilient nursery management through quantitative, data-driven methodologies. The integration of Internet of Things (IoT) sensor arrays, precision microfluidics, automated micropropagation bioreactors, tunable LED lighting, and artificial intelligence has enabled a paradigm shift toward Controlled Environment Agriculture. Five distinct architectural layers are examined: sensing/perception, connectivity/network, edge/cloud intelligence, actuation/control, and supervisory interface. Precision hydro-nurturing strategies demonstrate water consumption reductions from 12,000 L/ha to 6,800 L/ha through sensor-based IoT and predictive AI scheduling, alongside 30–70% reductions in fertilizer and pesticide applications. Automated Temporary Immersion Systems address scalability bottlenecks in micropropagation while preventing hyperhydricity and reducing labor costs. Spectral optimization through tunable LEDs reveals that red-blue-far-red combinations enhance root architecture and stress hardiness, while green supplementation achieves superior stem diameters (11.08 mm) and root lengths (49 cm). Predictive growth modeling integrating computer vision, functional-structural plant models, and digital twins enables non-destructive phenotyping and real-time stress forecasting. Economic barriers including high capital expenditure and digital literacy gaps are addressed through modular, open-source IoT pathways. Future horizons emphasize self-calibrating sensors, Explainable AI frameworks, and closed-loop stress pre-conditioning systems that prime seedling physiological pathways for enhanced field survival under changing climate conditions.

1. INTRODUCTION

Global agricultural and forestry sectors face unprecedented operational vulnerabilities driven by anthropogenic climate volatility, characterized by erratic precipitation patterns, unseasonal heatwaves, extended drought periods, and shifting pathogen pressure. Early-stage plant propagation represents the most environmentally vulnerable phase of the crop lifecycle. During this initial stage, seedling vigor, root system architecture, and anatomical establishment dictate subsequent field survival, stress tolerance, and ultimate economic yield (Grossnickle & MacDonald, 2018). Traditional nursery operations rely on open-field or basic shade-house facilities managed via empirical observation and fixed, calendar-based cultivation practices. These conventional approaches suffer from substantial water wastage, non-uniform growth, high sapling mortality from root-zone hypoxia or pathogens, and poor resource use efficiency (Landis et al., 1998).

To protect early-stage propagation from environmental risks, Controlled Environment Agriculture (CEA) and Smart Greenhouses (SGHs) have evolved into critical production frameworks. The convergence of Internet of Things (IoT) sensor arrays, precision microfluidics, automated micropropagation bioreactors, solid-state light-emitting diode (LED) lighting, and artificial intelligence (AI) has shifted nursery management toward a quantitative, data-driven methodology (Hosny et al., 2025). Beyond automated microclimate regulation, modern smart nurseries use predictive seedling growth modeling, high-throughput computer vision, and digital twins. These technologies allow operators to monitor physiological and morphological development non-destructively while dynamically forecasting crop trajectories and simulating stress responses (Singh et al., 2021).

This review provides a comprehensive synthesis of modern smart propagation technologies, cyber-physical IoT architectures, precision hydro-nurturing strategies, tunable LED photomorphogenesis, automated tissue culture systems, and predictive growth models. By examining the integration of physical hardware layers with biophysical computational engines, this paper highlights how smart nurseries cultivate physiological stress resilience in seedlings, optimize resource efficiency, and ensure robust field performance under changing climate conditions (Ariesen-Verschuur et al., 2022).




Figure 1: Comprehensive Workflow of Plant Propagation Pathways and Sequential Nursery Management Operations.

2. Architectural Frameworks and IoT Sensor Networks for Precision Nursery Infrastructure

The foundational engine of a climate-resilient smart nursery is a multi-tiered cyber-physical Internet of Things (IoT) framework that transforms continuous physical microclimatic and substrate inputs into automated agronomic decisions. Modern smart nursery architectures integrate five distinct operational layers: the Sensing/Perception Layer, the Connectivity/Network Layer, the Edge/Cloud Intelligence Layer, the Actuation/Hydraulic Control Layer, and the Supervisory Management Interface (Liakos et al., 2018).

The Sensing Layer continuously tracks microclimatic, substrate, and crop physiological indicators. Environmental arrays measure ambient air temperature, relative humidity, vapor pressure deficit (VPD), carbon dioxide (CO_2) concentration, barometric pressure, and photosynthetically active radiation (PAR) (Kamilaris & Prenafeta-Boldú, 2018). Substrate dynamics are tracked using Time Domain Reflectometry (TDR) probes, Frequency Domain Reflectometry (FDR) sensors, dielectric capacitance sensors (such as EC-5 probes), tensiometers, and electrical conductivity (EC) / pH sensors. Plant-centric sensing directly monitors crop stress via infrared canopy thermography to track the crop water stress index (CWSI), alongside stem dendrometers, sap flow sensors, leaf turgor pressure probes, and Normalized Difference Vegetation Index (NDVI) spectral sensors that quantify internal water status, vascular transport, and pigment density (Ariesen-Verschuur et al., 2022).

The Connectivity and Network Layer manages continuous data transmission across distributed nursery bays. Low-Power Wide-Area Networks (LPWAN) specifically Narrowband IoT (NB-IoT) and LoRaWAN alongside standard GSM and Wi-Fi protocols, provide reliable wireless sensor network (WSN) topologies with low energy consumption. NB-IoT architectures are well-suited for rural or large-scale commercial nurseries due to their deep signal penetration through greenhouse frames, reducing hardware implementation costs by 17.5% to 29.3% compared to traditional wired industrial systems (Wolfert et al., 2017).

Data processing is partitioned between Edge and Cloud Intelligence Layers. Local edge nodes handle time-critical safety loops such as immediate solenoid valve actuation during sudden substrate moisture drops ensuring uninterrupted operation during network outages. Concurrently, cloud infrastructure aggregates long-term time-series data, training deep learning growth models, executing Functional-Structural Plant Models (FSPMs), and updating predictive irrigation schedules (Ayaz et al., 2019). The Actuation Layer executes these cloud and edge decisions via relay interfaces that drive variable-speed pumps, nutrient dosing injectors, automated shade curtains, HVAC units, fogging systems, and dimmable LED arrays. Finally, the Supervisory Interface presents real-time analytics, yield forecasts, and system alerts to nursery managers (Dumroese et al., 2016).

Table 1. Architectural Layers and Technological Components of Smart Nursery Systems

Layer	Primary Hardware & Protocols	Agronomic Functions & Key Metrics	Primary Operational Benefits
Sensing & Perception	TDR/FDR probes, EC-5 sensors, canopy thermography, sap flow, leaf turgor, NDVI, ambient climate sensors	Volumetric water content (VWC), EC, pH, vapor pressure deficit (VPD), crop water stress index (CWSI)	Continuous detection of physiological plant stress and substrate nutrient status.
Connectivity & Network	NB-IoT, LoRaWAN, GSM links, Wi-Fi, LPWAN topologies	Data packet routing, duty-cycle scheduling, wireless mesh network maintenance	17.5–29.3% reduction in total hardware costs; robust long-range connectivity.

Edge & Cloud Intelligence	Microcontrollers (ESP32/Arduino), edge AI modules, cloud servers, time-series ML models	Sensor auto-calibration, 10–25% gain in moisture prediction accuracy, model updates	Low-latency edge execution, system operational resilience, deep trend analysis.
Actuation & Control	Solenoid valves, variable-frequency dosing pumps, PWM LED drivers, motor drives	Precision water delivery, fertigation dosing, dynamic spectral tuning, misting control	Automated execution of hydro-nurturing rules; removal of manual operational errors.
Supervisory Interface	Web dashboards, mobile applications, SCADA platforms, digital twin interfaces	Yield/growth trajectory tracking, alert management, manual overrides	Centralized operational control, parameter configuration, remote nursery monitoring.

3. Precision Hydro-Nurturing and Nutrient Management Strategies

Substrate moisture management in young seedlings directly influences root architecture, stomatal regulation, and vegetative growth. Traditional nursery irrigation typically operated via fixed timers or overhead impact sprinklers suffers from non-uniform distribution, root-zone saturation, nutrient leaching, and high incidence of fungal pathogens like *Pythium* and *Phytophthora*. Conversely, underwatering causes stomatal closure, suppresses net carbon assimilation, and stunts seedling growth (Lesk et al., 2016).

Smart hydro-nurturing replaces empirical watering schedules with closed-loop precision techniques. By linking real-time IoT substrate monitoring with automated drip delivery, water is applied strictly according to plant transpiration rates and substrate depletion. Comparative assessments across scheduling frameworks highlight significant water savings and growth improvements. Fixed-schedule manual irrigation consumes approximately 12,000 L/ha, with water wastage exceeding 20%. Sensor-based IoT systems reduce consumption to 7,200 L/ha while reducing wastage by 15% and boosting overall crop yield. Combining real-time IoT monitoring with predictive AI algorithms further optimizes delivery to 6,800 L/ha (IPCC, 2022).

The choice of physical water delivery technology also impacts root-zone dynamics. Precision drip systems reduce water usage by 30% to 70%, chemical fertilizer requirements by 30% to 60%, and pesticide applications by 40% to 50% relative to surface flooding or overhead watering, while increasing overall biomass and crop yield by 20% to 40%. Soil water controllers interfaced with dielectric probes reduce water usage by up to 61% compared to traditional evapotranspiration-based scheduling by responding directly to physical substrate water retention dynamics (Davis & Jacobs, 2005).

Table 2. Comparative Assessment of Nursery Irrigation Scheduling Frameworks and Resource Conservation Metrics

Irrigation Scheduling Framework	Estimated Water Consumption (L/ha)	Relative Water Savings (%)	Water Wastage Reduction (%)	Agronomic & Crop Vigor Impact
Manual (Fixed Schedule)	12,000	Baseline (0%)	Baseline	High risk of substrate hypoxia, nutrient leaching, and non-uniform seedling establishment.
Volume-Based Scheduling	10,500	12.5%	+2% efficiency gain over baseline	Moderate reduction in excess run-off; fails to adapt to rapid VPD fluctuations.

Sensor-Based IoT Scheduling	7,200	40.0%	15% reduction in total wastage	Maintains steady volumetric moisture, improves root zone aeration, drives biomass gains.
Real-Time IoT + AI Predictive Models	6,800	43.3%	>15% reduction in total wastage	Optimizes water use efficiency (WUE), stomatal conductance, and stress mitigation.

Substrate volumetric water content (VWC) stability also dictates nutrient solubility and mineral uptake. Controlled trial evaluations comparing IoT-managed greenhouses to traditional farming setups demonstrate that real-time sensor control stabilizes gravimetric water content and prevents sudden substrate pH fluctuations. This stable moisture regime significantly increases plant retention and root absorption of key micronutrients and secondary macronutrients, including iron (Fe), manganese (Mn), calcium (Ca), magnesium (Mg), and zinc (Zn). In a 120-day comparative growth study, tomato plants managed under real-time IoT greenhouse controls achieved significantly greater height compared to open control plots (mean height difference = 0.3564 m, $p = 0.001$, 95% CI: [0.1806, 0.5323]). This enhanced growth stems from steady nutrient availability under controlled substrate electrical conductivity, preventing nutrient immobilization caused by substrate drying or flushing losses from over-irrigation (Shamshiri et al., 2018).

4. Advanced Micropropagation and Automated Bioreactor Systems

For vegetatively propagated crops, elite horticultural cultivars, and forestry species, micropropagation provides rapid clonal expansion, year-round production, and pathogen-free starting material (Kozai et al., 2019). Regulatory phase-outs of chemical soil fumigants like methyl bromide, alongside compounding pressures from soil-borne pathogens such as *Colletotrichum gloeosporioides*, *Phytophthora cactorum*, and *Verticillium dahliae*, have accelerated the shift from field nurseries to controlled micropropagation platforms. Traditional tissue culture on semi-solid agar media, however, remains labor-intensive, subject to vessel handling constraints, restricted by poor vessel gas exchange, and prone to physiological disorders such as hyperhydricity (Paek et al., 2005).

To address these scalability bottlenecks, automated liquid culture technology specifically Temporary Immersion Systems (TIS) like RITA bioreactors and twin-bottle systems has become the modern standard for semi-automated micropropagation. TIS operates by periodically surging liquid nutrient media into the tissue chamber to submerge explants for set durations, followed by complete medium pneumatic drainage (Paradiso & Proietti, 2022).

This periodic immersion cycle provides key physiological benefits:

- Pneumatic venting during the dry phase refreshes internal vessel oxygen and carbon dioxide levels, stimulating photosynthetic capacity and shoot differentiation.
- Explants are left with a thin liquid film covering their surfaces after medium drainage, facilitating nutrient uptake without inducing cellular hypoxia.
- Avoiding continuous submersion prevents tissue hyperhydricity, producing plants with functional cuticular waxes and responsive stomata.
- TIS eliminates manual subculturing onto solid agar media, lowering labor costs and facilitating integration with automated liquid-handling systems (Klerkx et al., 2019).

The transition of micropropagated plantlets from *in vitro* vessel environments to *ex vitro* nursery substrates represents a vulnerable acclimatization phase. In vitro plantlets typically possess poorly developed cuticular wax layers, non-functional stomatal complexes, and weak vascular connections between roots and shoots. Modern smart nurseries deploy specific physiological hardening strategies to maximize ex vitro survival.

Automated misting and fogging bays continuously modulate relative humidity, stepping it down gradually from 95% to ambient levels over 10–14 days to stimulate epicuticular wax synthesis and stomatal functionality (Grossnickle, 2012). Simultaneously, bio-inoculation with Arbuscular Mycorrhizal Fungi (AMF) during transplanting expands root absorptive surface area, improves phosphorus uptake, and strengthens systemic abiotic stress tolerance. Furthermore, foliar applications of natural elicitors like chitosan at concentrations of 0.15% to 0.20% increase stem diameter, leaf area, structural biomass, and chlorophyll content while upregulating endogenous defense pathways (Grossnickle & MacDonald, 2018).




Figure 2: Biotechnological Strategy for Developing Climate-Resilient Apomictic Crop Varieties via Trait Identification, Gene Editing, and Clonal Seed Production.

5. Photomorphogenic Optimization via Tunable LED Lighting Regimes

Light quality (spectral wavelength composition), quantity (Photosynthetic Photon Flux Density, PPFD), and photoperiod function as critical environmental cues that direct plant photomorphogenesis. Solid-state LED technology enables precise spectral tuning, allowing nursery operators to manipulate specific wavelengths to induce targeted anatomical and structural traits (van Iersel et al., 2016).

Plant light perception is mediated by dedicated photoreceptors: phytochromes absorb red and far-red spectra, cryptochromes and phototropins respond to blue and UV-A wavelengths, and zeitlupe family proteins manage circadian rhythms. Manipulating these spectral bands triggers distinct developmental pathways (Smith, 2020).

Red light ($\lambda \approx 660$ nm) aligns with the absorption peaks of chlorophyll a and b, converting inactive phytochrome P_r into active P_{fr}. This transition drives photosynthetic carbon assimilation, leaf blade expansion, and overall biomass accumulation. However, exposure to monochromatic red-light causes "red-light syndrome," characterized by stomatal dysfunction, stem etiolation, and reduced photosynthetic efficiency (Quail, 2002).

Blue light ($\lambda \approx 450$ nm) activates cryptochromes, suppressing internode elongation and producing compact, sturdy plant structures with thicker leaves and higher stomatal densities. Higher blue light proportions upregulate chlorophyll synthesis, secondary metabolite production, and structural root development. Combining red and blue light (e.g., a 10:1 R:B ratio) produces optimal morphological outcomes in young plantlets, yielding higher leaf counts, leaf area expansion, root counts, root length, and total dry biomass compared to monochromatic light (Casal et al., 2014).

Far-red light ($\lambda \approx 730$ nm) converts active P_{fr} back to inactive P_r, lowering the phytochrome photostationary state (PSS) and simulating shade avoidance responses. Low Red to Far-Red (R:FR) ratios stimulate petiole and stem elongation, apical dominance, and leaf surface area expansion. While excessive far-red can cause unwanted stem stretching, moderate far-red supplementation boosts whole-plant net assimilation by expanding the light-capturing leaf canopy (Klose et al., 2015). Multi-spectral combinations that incorporate far-red into red/blue bases (such as 2 Parts Red : 1 Part Blue : 1 Part Far-Red) enhance root system architecture by increasing total root volume, length, and lateral root branching without triggering excessive shoot elongation. This balanced architecture improves root-to-shoot biomass ratios, increasing post-transplant survival in field environments. Similarly, adding green light (such as Treatment L4, an RGB blend) achieves optimal stem diameters (11.08 mm), extended root length (49 cm), and superior overall plant fresh weight (Sanchez et al., 2020).

Table 3. Morphological and Physiological Responses of Seedlings to Specific LED Spectral Compositions

LED Spectral Composition	Primary Photoreceptors & Signaling Pathways	Dominant Morphological Response	Key Physiological & Biochemical Effects	Ideal Nursery Application Phase
Monochromatic Red (100% R)	Phytochrome conversion ($P_r \rightarrow P_{fr}$)	Pronounced stem elongation, hypocotyl stretching, reduced branching	High photosynthetic drive; risk of stomatal dysfunction ("red-light syndrome")	Early seed germination and shoot emergence.
Monochromatic Blue (100% B)	Cryptochrome / Phototropin activation	Dwarf growth habit, internode inhibition, compact structure, thick leaves	Elevated stomatal conductance, enhanced chlorophyll synthesis, metabolite accumulation	Vegetative compacting; preventing leggy growth.
Red + Blue (RB Ratio 10:1)	Dual Phytochrome & Cryptochrome activation	Maximum leaf surface area, high node/leaf counts, robust shoot and root numbers	Highest chlorophyll a/b accumulation, optimal net photosynthesis, high dry mass	Standard micropropagation shoot multiplication and plug production.
Red + Blue + Far-Red (2R : 1B : 1FR)	Phytochrome photostationary state (PSS) modulation	Enhanced root branching, increased root length/volume,	Balanced root-to-shoot biomass allocation,	Pre-transplant root hardening and acclimatization.

		controlled shoot height	improved systemic stress hardiness	
Red + Blue + Green (RBG / Treatment L4)	Balanced RGB photoreceptor activation	Superior stem diameter (11.08 mm), broad leaves, long roots (49 cm) and stolons	GA3 hormonal regulation, high net assimilation rates, superior plant fresh weight	General transplant development and aeroponic growth.

Spectral composition also controls morphological development by modulating endogenous hormone networks. Red-dominant spectra stimulate gibberellin (GA_3) biosynthesis, promoting cell division and internode stretching. Low R:FR light ratios accelerate polar auxin transport from shoot tips to root zones while downregulating jasmonic acid (JA) and salicylic acid (SA) defense pathways. This physiological tradeoff reallocates metabolic resources toward rapid physical elongation rather than immune defense, making spectral management essential to prevent increased disease susceptibility in commercial nurseries (Wang et al., 2024).

6. Predictive Growth Modeling, Computer Vision, and Digital Twin Integration

Modern smart nurseries increasingly complement physical hardware with computational modeling, replacing manual visual inspection with non-destructive, continuous phenotyping. Combining high-throughput computer vision, machine learning (ML), functional-structural plant models (FSPMs), and digital twin platforms allows operators to evaluate seedling vigor and dynamically adjust growing environments (Li et al., 2020; Murphy et al., 2024; Rahman et al., 2024).

Non-destructive phenotyping relies on RGB, Multispectral Imaging (MSI), and Hyperspectral Imaging (HSI) vision sensors. RGB cameras capture structural metrics, including projected canopy area, leaf count, stem height, and color indices. Multispectral and hyperspectral cameras detect subtle variations in chlorophyll absorption, water band reflections, and anthocyanin accumulation. These spectral signatures allow automated vision systems to identify nutrient deficiencies, water stress, or early disease onset before visible symptoms appear (Kuska et al., 2015).

Deep learning models process time-series image sequences to forecast future seedling growth trajectories. Generative deep learning networks and convolutional regression models process historical image sequences to synthesize high-resolution predicted images of plant canopy structures at future time steps (image). These models automatically extract up to 37 distinct phenotypic traits, including leaf blade perimeter, plant height, total projected leaf area, internode distance, and canopy density. Self-supervised models map these visual inputs onto dynamic growth trajectories, enabling systems to predict market readiness dates, detect growth stunting early, and auto-correct fertigation or lighting treatments (Sakurai et al., 2025).

While purely statistical machine learning models excel at image pattern recognition, they lack underlying biophysical mechanics. Functional-Structural Plant Models (FSPMs) resolve this limitation by combining 3D structural architecture modeling with biophysical process equations for photosynthesis, transpiration, carbon allocation, and hormonal signaling (Vos et al., 2010).

Integrating FSPMs with continuous IoT sensor feeds creates a real-time Digital Twin of the nursery crop. A Digital Twin forms a bi-directional virtual-physical feedback system. High-throughput vision sensors and root-zone IoT nodes stream operational data into the digital twin platform. The underlying 3D FSPM updates its physiological state variables in real time, calculating carbon allocation, biomass accumulation, and transpiration rates under ambient conditions. Using these simulations, the Digital Twin forecasts growth under potential microclimatic stress

scenarios, such as heat spikes or dry spells. It then transmits optimized actuation commands back to edge controllers to adjust light spectra, irrigation frequency, or shading systems, ensuring seedlings remain on target growth trajectories (Sievänen et al., 2014).

7. Economic Feasibility, Technological Bottlenecks, and Scalability Pathways

Despite the agronomic and resource efficiency benefits of smart propagation systems, commercial adoption faces economic and technical challenges. High initial capital expenditure (CAPEX) for advanced sensor networks, CEA infrastructure, automated TIS bioreactors, multi-channel LED fixtures, and cloud software platforms present a barrier for small- and medium-scale commercial nurseries. Additionally, operating advanced software platforms, deep learning vision pipelines, and automated fertigation systems requires specialized technical expertise, creating a digital literacy gap among traditional nursery staff (Singh et al., 2022).

To improve technological accessibility, recent engineering efforts focus on open-source, modular, and low-cost IoT systems. Standardizing communication architectures around Narrowband IoT (NB-IoT) reduces system hardware costs by 17.5% to 29.3% compared to conventional industrial automation networks while maintaining reliable data collection and control accuracy. Open-source microcontrollers, such as ESP32 or Arduino platforms, paired with standard dielectric probes, provide accessible entry points for precision hydro-nurturing. Modular adoption pathways where facilities start with automated drip irrigation before adding spectral LED arrays and phenotyping platforms offer a practical approach for progressive technological integration (Sagan et al., 2019).

8. Strategic Roadmap and Future Horizons

To advance climate-resilient nursery management, ongoing research focuses on three strategic development priorities:

First, sensor technology is moving toward self-calibrating and auto-correcting IoT nodes. Substrate-embedded sensors often experience signal drift and calibration loss due to biofouling or salt accumulation. Sensor nodes featuring embedded machine learning for real-time drift detection, cross-validation, and auto-calibration against reference arrays will lower maintenance demands and extend operational lifespans (Murphy et al., 2024).

Second, decision-support tools are adopting Trustworthy and Explainable AI (XAI) frameworks. Black-box deep learning models often face adoption resistance from nursery managers who require transparent rationale for automated microclimate interventions. Future predictive growth engines must combine data-driven neural networks with biophysical plant physiology principles, generating interpretable, actionable agronomic recommendations (Mahlein et al., 2018).

Third, nurseries are moving toward closed-loop stress pre-conditioning systems. Future facilities will leverage real-time digital twins to apply controlled, non-lethal micro-stressors such as temporary vapor pressure deficit spikes, dynamic far-red light pulses, or mild water deficits. These controlled interventions prime seedling physiological pathways, upregulating protective heat-shock proteins, strengthening root architecture, and thickening cuticular wax layers. This stress hardening ensures high seedling survival and rapid establishment when transplanted into challenging field environments (Runions et al., 2017).

Conclusion

This review demonstrates that climate-resilient nursery management through smart propagation technologies represents a fundamental paradigm shift from traditional empirical practices to quantitative, data-driven methodologies. The integration of multi-tiered IoT architectures, precision hydro-nurturing strategies, automated micropropagation systems, tunable LED photomorphogenesis, and predictive growth modeling creates a cohesive cyber-physical framework capable of cultivating physiological stress resilience in seedlings while optimizing

resource efficiency. The five-layer architectural framework encompassing sensing, connectivity, edge/cloud intelligence, actuation, and supervisory interfaces provides the foundational infrastructure for continuous monitoring and automated decision-making. Precision irrigation scheduling frameworks achieve water savings of 40–43.3% compared to manual fixed schedules, with sensor-based IoT systems reducing consumption from 12,000 L/ha to 7,200 L/ha and AI-predictive models further optimizing delivery to 6,800 L/ha. These reductions are accompanied by improved root zone aeration, stabilized nutrient availability, and enhanced crop vigor. Automated Temporary Immersion Systems overcome scalability bottlenecks inherent in traditional semi-solid agar culture, delivering physiological benefits including improved vessel gas exchange, prevention of hyperhydricity, and facilitation of automated liquid-handling integration. Spectral optimization through tunable LEDs enables precise manipulation of photomorphogenic pathways, with multi-spectral combinations incorporating far-red and green wavelengths producing superior root architecture, stem diameter, and biomass allocation. These morphological advantages translate directly to improved post-transplant survival in challenging field environments. Predictive growth modeling through computer vision, deep learning, and functional-structural plant models enables non-destructive, continuous phenotyping capable of forecasting growth trajectories and detecting stress before visible symptoms manifest. Digital twin integration creates bi-directional virtual-physical feedback systems that simulate stress scenarios and transmit optimized actuation commands to maintain seedlings on target growth trajectories. Despite these advances, commercial adoption faces barriers including high capital expenditure, specialized technical expertise requirements, and digital literacy gaps. Modular, open-source IoT systems and standardized communication architectures reduce hardware costs by 17.5–29.3%, offering accessible entry points for progressive technological integration. Future development priorities include self-calibrating sensor nodes with embedded machine learning for drift detection, Trustworthy and Explainable AI frameworks that combine data-driven neural networks with biophysical plant physiology principles, and closed-loop stress pre-conditioning systems that apply controlled non-lethal micro-stressors to prime protective physiological pathways. Collectively, these innovations position smart nurseries as critical infrastructure for ensuring robust field performance and food security under accelerating climate change.

REFERENCES

Ariesen-Verschuur, N., Verdouw, C., & Tekinerdogan, B. (2022). Digital twins in greenhouse horticulture: A review. *Computers and Electronics in Agriculture, 199*, 107183. https://doi.org/10.1016/j.compag.2022.107183

Ayaz, M., Ammad-Uddin, M., Sharif, Z., Mansour, A., & Aggoune, E.-H. M. (2019). Internet-of-Things (IoT)-based smart agriculture: Toward making the fields talk. *IEEE Access, 7*, 129551–129583. https://doi.org/10.1109/ACCESS.2019.2932609

Casal, J. J., Candia, A. N., & Sellaro, R. J. J. E. B. (2014). Light perception and signalling by phytochrome A. *Journal of Experimental Botany*, *65*(11), 2835-2845.

Davis, A. S., & Jacobs, D. F. (2005). Quantifying root system quality of nursery seedlings and relationship to outplanting performance. *New Forests, 30*, 295–311. https://doi.org/10.1007/s11056-005-7480-5

Dumroese, R. K., Landis, T. D., Pinto, J. R., & Haase, D. L. (2016). *Using container seedlings to restore drylands: A practical guide*. U.S. Department of Agriculture, Forest Service.

Grossnickle, S. C. (2012). Why seedlings survive: Influence of plant attributes. *New Forests, 43*, 711–738. https://doi.org/10.1007/s11056-012-9336-6

Grossnickle, S. C., & MacDonald, J. E. (2018). Seedling quality: History, application, and plant attributes. *Forests, 9*(5), 283. https://doi.org/10.3390/f9050283

Hosny, K. M., El-Hady, W. M., & Samy, F. M. (2025). Technologies, protocols, and applications of Internet of Things in greenhouse farming: A survey of recent advances. *Information Processing in Agriculture, 12*(1), 91–111. https://doi.org/10.1016/j.inpa.2024.04.002

Intergovernmental Panel on Climate Change. (2022). *Climate change 2022: Impacts, adaptation and vulnerability*. Cambridge University Press. https://doi.org/10.1017/9781009325844

Kamilaris, A., & Prenafeta-Boldú, F. X. (2018). Deep learning in agriculture: A survey. *Computers and Electronics in Agriculture, 147*, 70–90. https://doi.org/10.1016/j.compag.2018.02.016

Klerkx, L., Jakku, E., & Labarthe, P. (2019). A review of social science on digital agriculture, smart farming and agriculture 4.0: New contributions and a future research agenda. *NJAS: Wageningen Journal of Life Sciences, 90–91*, 100315. https://doi.org/10.1016/j.njas.2019.100315

Klose, C., Viczián, A., Kircher, S., Schäfer, E., & Nagy, F. (2015). Molecular mechanisms for mediating light-dependent nucleo/cytoplasmic partitioning of phytochrome photoreceptors. *New Phytologist*, *206*(3), 965-971.

Kozai, T., Niu, G., & Takagaki, M. (Eds.). (2019). *Plant factory: An indoor vertical farming system for efficient quality food production* (2nd ed.). Academic Press.

Kuska, M. T., Behmann, J., Mahlein, A.-K., & Walter, A. (2015). Hyperspectral imaging for plant phenotyping and stress detection. *Plant Methods, 11*, 1–10.

Landis, T. D., Tinus, R. W., McDonald, S. E., & Barnett, J. P. (1998). *The container tree nursery manual: Seedling propagation*. U.S. Department of Agriculture, Forest Service.

Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature, 529*, 84–87. https://doi.org/10.1038/nature16467

Li, Z., Guo, R., Li, M., Chen, Y., & Li, G. (2020). A review of computer vision technologies for plant phenotyping. *Computers and Electronics in Agriculture, 176*, 105672. https://doi.org/10.1016/j.compag.2020.105672

Liakos, K. G., Busato, P., Moshou, D., Pearson, S., & Bochtis, D. (2018). Machine learning in agriculture: A review. *Sensors, 18*(8), 2674. https://doi.org/10.3390/s18082674

Mahlein, A.-K., Kuska, M. T., Behmann, J., Polder, G., & Walter, A. (2018). Hyperspectral sensors and imaging technologies in phytopathology: State of the art. *Annual Review of Phytopathology, 56*, 535–558. https://doi.org/10.1146/annurev-phyto-080417-050100

Murphy, K. M., Ludwig, E., Gutierrez, J., & Gehan, M. A. (2024). Deep learning in image-based plant phenotyping. *Annual Review of Plant Biology, 75*, 771–795. https://doi.org/10.1146/annurev-arplant-070523-042828

Paek, K.-Y., Hahn, E.-J., & Son, S.-H. (2005). Application of bioreactors for large-scale micropropagation systems of plants. *In Vitro Cellular & Developmental Biology Plant, 41*, 122–128. https://doi.org/10.1079/IVP2004600

Paradiso, R., & Proietti, S. (2022). Light-quality manipulation to control plant growth and photomorphogenesis in greenhouse horticulture: The state of the art and the opportunities of modern LED systems. *Journal of Plant Growth Regulation, 41*, 742–780. https://doi.org/10.1007/s00344-021-10337-y

Quail, P. H. (2002). Photosensory perception and signalling in plant cells: new paradigms?. *Current opinion in cell biology*, *14*(2), 180-188.

Rahman, H., Shah, U. M., Riaz, S. M., Kifayat, K., Moqurrab, S. A., & Yoo, J. (2024). Digital twin framework for smart greenhouse management using next-gen mobile networks and machine learning. *Future Generation Computer Systems, 156*, 285–300. https://doi.org/10.1016/j.future.2024.03.023

Runions, A., Tsiantis, M., & Prusinkiewicz, P. (2017). A common developmental program can produce diverse leaf shapes. *New Phytologist, 216*(2), 401–418.

Sagan, V., Maimaitijiang, M., Bhadra, S., Maimaitiyiming, M., Brown, D. R., Sidike, P., & Fritschi, F. B. (2019). Field-scale crop stress identification by multi-temporal deep learning. *Remote Sensing, 11*(13), 1–21.

Sanchez, S. E., Rugnone, M. L., & Kay, S. A. (2020). Light perception: a matter of time. *Molecular plant*, *13*(3), 363-385.

Shamshiri, R. R., Kalantari, F., Ting, K. C., Thorp, K. R., Hameed, I. A., Weltzien, C., Ahmad, D., & Shad, Z. M. (2018). Advances in greenhouse automation and controlled environment agriculture: A transition to plant factories and urban agriculture. *International Journal of Agricultural and Biological Engineering, 11*(1), 1–22. https://doi.org/10.25165/IJABE.V11I1.3210

Sievänen, R., Godin, C., DeJong, T. M., & Nikinmaa, E. (2014). Functional–structural plant models: A growing paradigm for plant studies. *Annals of Botany, 114*(4), 599–603. https://doi.org/10.1093/aob/mcu175

Singh, A., Ganapathysubramanian, B., Singh, A. K., & Sarkar, S. (2022). Machine learning for high-throughput stress phenotyping in plants. *Trends in Plant Science, 27*(3), 301–315.

Singh, V., Sharma, N., & Singh, S. (2021). A review of imaging techniques for plant disease detection and diagnosis. *Computers and Electronics in Agriculture, 184*, 106089.

Smith, H. (2000). Phytochromes and light signal perception by plants an emerging synthesis. *Nature*, *407*(6804), 585-591.

van Iersel, M. W., Gianino, D., & Van Pelt, J. (2016). Supplemental lighting in controlled environment agriculture. *HortScience, 51*, 121–128.

Vos, J., Evers, J. B., Buck-Sorlin, G. H., Andrieu, B., Chelle, M., & de Visser, P. H. B. (2010). Functional–structural plant modelling: A new versatile tool in crop science. *Journal of Experimental Botany, 61*(8), 2101–2115. https://doi.org/10.1093/jxb/erp345

Wang, Q., Bian, Z., Wang, S., Zhao, Y., Zhan, X., & Yang, Q. (2024). Far-red light inhibits soybean biomass and yield by modulating plant photosynthesis. *Agronomy*, *14*(11), 2684.

Wolfert, S., Ge, L., Verdouw, C., & Bogaardt, M.-J. (2017). Big data in smart farming A review. *Agricultural Systems, 153*, 69–80. https://doi.org/10.1016/j.agsy.2017.01.003

IEER
Excellence in Education & Research